



The **Economic benefits** of the **Natura 2000** Network



Cover photo: istockphoto

The cliffs of Moher, Ireland, is a Natura 2000 site, a Special Protection Area for seabirds and a major tourist destination.

This document arises from a contract for the European Commission: Estimating the Overall Economic Value of the Benefits provided by the Natura 2000 Network. Final Synthesis Report to the European Commission, DG Environment on Contract 07.0307/2010/581178/SER/B3, Brussels 2011. It was prepared by Institute for European Policy (IEEP): P ten Brink, S Bassi, T Badura, S Gantioler, M Kettunen, L Mazza, K Hart together with GHK: M Rayment, M Pieterse, E Daly, Ecologic Institute: H Gerdes, M Lago, S Lang, Metroeconomica: A Markandya, P Nunes, H Ding and EFTEC: R Tinch, I Dickie.

Acknowledgements

The contribution of Dr. Mike Christie, Stephen Nicol and Dr. Unai Pascual, as reviewers of the study is acknowledged, as is the valuable inputs of Johan Lammerant and his study team from Arcadis Belgium, EFTEC and ECNC for the project 'Recognising Natura 2000 Benefits And Demonstrating The Economic Benefits Of Conservation Measures', and Blandine Chenot and her study team from BIO Intelligence Service, Ecotrans, OÄR and Dunira Strategy for another project 'Estimating the economic value of the benefits provided by the tourism/recreation and employment supported by Natura 2000'.

***Europe Direct is a service to help you find answers
to your questions about the European Union***

Freephone number (*):

00 800 6 7 8 9 10 11

(*) Certain mobile telephone operators do not allow access to 00 800 numbers or these calls may be billed.

More information on the European Union is available on the Internet (<http://europa.eu>).

Cataloguing data can be found at the end of this publication.

Luxembourg: Publications Office of the European Union, 2013

ISBN 978-92-79-27588-3

doi:10.2779/41957

© European Union, 2013

Reproduction is authorised provided the source is acknowledged.

Printed in Belgium

PRINTED ON RECYCLED PAPER THAT HAS BEEN AWARDED THE EU ECO-LABEL FOR GRAPHIC PAPER

(WWW.ECOLABEL.EU)

The **Economic benefits** of the Natura 2000 Network

Synthesis Report



Foreword



The EU's economic prosperity and well-being is underpinned by its natural capital, which includes natural ecosystems that provide essential goods and services. Knowledge concerning the flow of benefits from ecosystems has grown considerably in recent years, encouraged by the work on the economics of ecosystems and biodiversity (TEEB) and the earlier Millennium Ecosystem Assessment (MA). Building on this approach, this study provides a first evaluation of the benefits of the Natura 2000 network, the EU network of area of high biodiversity value.

It clearly demonstrates that Natura 2000 has a key role to play in protecting and enhancing our natural capital. In addition to safeguarding nature's intrinsic value, investing in Natura 2000 provides multiple benefits to society and the economy at the local, regional, national and EU level.

The network is a major store of carbon rich habitats and has an important role to play in responding to the challenges we face from climate change, both through mitigation and adaptation. It also delivers other socio-economic benefits such as maintaining water flow and quality, conserving natural pollinators, preserving landscape and amenity values, and supporting tourism and recreation.

According to this study, the benefits that flow from Natura 2000 are of the order of €200 to 300 billion/year. It is estimated that there are between 1.2 to 2.2 billion visitor days to Natura 2000 sites each year, generating recreational benefits worth between €5 and €9 billion per annum. Therefore, investing in Natura 2000 makes sense and is directly relevant to Europe 2020 objectives of growth and employment as it can be a motor for the local and regional economy.

Natura 2000, as a key element of Green Infrastructure, also helps to safeguard the flow of ecosystem services that are otherwise at risk of degradation. Investment in management and restoration measures can increase the provision of a range of the services, from the scientific valuation of sites to flood control and water purification as the conservation status of the sites improves. The strong legal protection that applies to Natura 2000 also has an added benefit, providing long-term security to any financial investments to safeguard the sites it contains and the benefits they deliver.

Securing the necessary support for Natura 2000 in these difficult economic times represents a significant challenge so it is vital to communicate the benefits provided by the network. Nature needs our help, but it will pay us back many times over. Everyone has a role to play in making Natura 2000 a success – be they public authorities, private landowners and users, developers, conservation NGOs, scientific experts, local communities. This is the legacy of partnership that we should aim to leave for future generations.

Janez Potočnik

European Commissioner for the Environment

Acronyms

CBD	Convention on Biological Diversity
GHG	Greenhouse Gas
CFP	Common Fisheries Policy
COPI	Cost of Policy Inaction
CVM	Contingent Valuation (Method)
EC	European Communities
EEA	European Environment Agency
ESS	Ecosystem Service
EU	European Union
FCS	Favourable Conservation Status
FTE	Full time Employment
GDP	Gross Domestic Product
GIS	Geographical Information System
HNV	High Nature Value
IEEP	Institute for European Environmental Policy
IUU	illegal, unreported and unregulated
JRC	Joint Research Centre
MA	Millennium Ecosystem Assessment
MPA	Marine Protected Area
MSY	Maximum Sustainable Yield
PA	Protected Area
PES	Payment For Ecosystem Service(s)
SAC	Special Areas of Conservation
SCI	Sites of Community Importance
SPA	Special Protection Areas
TEEB	The Economics of Ecosystems and Biodiversity
TEV	Total Economic Value
UK	United Kingdom
UN	United Nations
UNEP	United Nations Environment Programme
WHO	World Health Organization
WTP	Willingness-To-Pay

Table of content

Part A: Aims and approach	7
I) Introduction: The Natura 2000 network and its benefits	7
II) Methodology for assessing EU wide benefits of Natura 2000	11
III) The total Value of the Natura 2000 network – a first assessment	17
Part B: Deriving an aggregate total value of services from natura 2000	17
IV) Overview of Ecosystem Services	23
Part C: The value of different Ecosystem services from Natura 2000	23
V) Natura 2000's fundamental role in climate mitigation – The Carbon storage and sequestration benefits of Natura 2000	25
VI) Natura 2000 as a tool for security: Natural hazards benefits and climate adaptation	28
VII) Natura 2000 as a motor of the economy/oil of society – The tourism and recreation benefits	30
VIII) Natura 2000 and 'free' resources for/value for money in the economy and society: Water purification and supply benefits (and waste)	32
IX) Natura 2000 and food: Marine protected areas and fish, and terrestrial protected areas, pollination and agriculture	35
X) Natural 2000 and our health, identity and learning	40
Part D: Realising the Benefits of Natura 2000	43
XI) Realising the Benefits: restoration and conservation for biodiversity and co-benefits.	43
Part E: Summary of results and outlook	49
XII) Summary of results	49
XIII) Outlook: improving the knowledge base for the economic valuation of ecosystem services delivered by Natura 2000	56
References	62
Annex I: Glossary of terms	66
Annex II: Overview of existing valuation site-based studies	68
Anex III: Examples of ecosystem services	71

Part A:

Aims and approach

I) Introduction: The Natura 2000 network and its benefits

The primary focus of the Natura 2000 protected area network is the conservation of biodiversity, the EU's unique and endangered ecosystems, species, gene pool and habitats. There has been an increasing, complementary interest in and recognition of the socio-economic benefits of biodiversity in general (MA, 2005; TEEB 2010, 2011) and from protected areas specifically (Kettunen et al 2009 & 2011, Stolton et al 2010, Gantioler 2010, Kettunen et al 2011) over the last decade.

In addition to its biodiversity benefits, the Natura 2000 network provides a range of benefits to society and the economy via the flow of ecosystem services (provisioning, regulating, cultural and supporting services). These support policy objectives beyond biodiversity, including climate change mitigation and adaptation, water quality and provision, food provision, jobs and livelihoods, cost savings, science and education, health and security, social cohesion and identity.

The recognition and demonstration of the wider socio-economic benefits of Natura 2000 can influence stakeholder attitudes and support for the Natura 2000 network, attract funding for conservation measures and other investment in and around sites, inform land-use (change) decisions, and help in the integration of protected areas in regional development planning and practice.

This report presents the results of a study to develop a methodological framework for assessing the overall economic value of the benefits provided by the Natura 2000 network, carrying out a first assessment of the value of the Natura 2000 network, and an outlook for future assessments to support the awareness of the economic co-benefits of the Natura 2000 sites and network.

In order to estimate the value of the network, the 'ecosystem services' framework has been adopted within this study, building on the Millennium Ecosystem Assessment (MA, 2005) and The Economics of Ecosystems and Biodiversity (TEEB, 2010 and 2011) – see the Glossary in Annex I for definitions and Chapter III and the full technical report of the study for wider discussion of the methodological framework.

Context: The Natura 2000 network, its coverage, rationale and benefits

The EU has a well-developed biodiversity conservation policy framework, which has been built up in response to international initiatives such as the Convention on Biological Diversity (CBD) and Bern Convention, and successive EU Environmental Action Programmes. In 2011 the European Commission adopted an ambitious new strategy¹ which provides the strategic framework for action to halt and reverse the loss of biodiversity and ecosystem services in the EU by 2020. At the heart of the EU's biodiversity conservation policy framework are the Birds Directive² and Habitats Directive³, which form the main legal framework for the protection of nature and biodiversity in the EU.

To achieve their objectives both Directives require two main types of activities. Firstly, the designation, implementation and management of sites that are particularly important for conserving and restoring EU biodiversity, and secondly, the strict protection of listed species as well as their breeding sites and resting places, wherever they occur. The establishment, protection and management of a coherent network of areas – known as the Natura 2000 network – is designed to protect the habitats and species targeted by the Directives.

¹ Communication from the Commission: Our life insurance, our natural capital: an EU biodiversity strategy to 2020 (COM(2011) 244)

² Council Directive 2009/147/EC on the conservation of wild birds (Birds Directive) adopted in 1979

³ Council Directive 92/43/EEC on the conservation of natural habitats and of wild fauna and flora (Habitats Directive) adopted in 1992

Figure 1: Natura 2000 areas

NATURA 2000: Birds and Habitats Directives

NATURA 2000

- Birds Directive sites (SPA)
- Habitats Directive sites (pSCI, SCI, SAC)
- Sites - or parts of sites - belonging to both directives

European Environment Agency




Source: Natura 2000- DGENV, compiled from databases from Member States.

Sources background map: EuroGlobalMap/Eurographics and DG ESTAT.

Validity of Natura 2000 data for Europe. Updated: August 2011.

Source: <http://www.eea.europa.eu/data-and-maps/figures/natura-2000-birds-and-habitat-directives-3>

Figure 2: The conservation status of habitats in the EU's biogeographic regions

The pie charts indicate for each biogeographical region the proportions of the conservation status assessments that are favourable, unfavourable and unknown.

Sources: ETC/BD, 2008; SEB1 2010 Indicator 05

The network is comprised of 26,400 sites and covers almost 18% of the EU territory. It includes terrestrial Sites of Community Importance (SCIs), with an area of 59 million ha (0.59 million km²), and terrestrial Special Protection Areas (SPAs) with an area of 52 million ha (0.52 million km²).

It also includes a growing marine protected area (MPA) network – now at 21.7 million ha⁴: 12 million ha² classify as SPAs and 18 million ha² as SCIs (note there is a significant number of sites that are both SCI and SPAs). The network is a core element of the wider EU green infrastructure, which together form a great part of our living natural capital.

The primary focus of the Natura 2000 protected area (PA) network is the conservation of the unique and endangered biodiversity in Europe; this includes rare habitats (e.g. cold water coral reefs), species (from keystone species to iconic charismatic species such as the Iberian Lynx) and genetic diversity (e.g. number of endemic species).

The benefits of protected areas to people, society and the economy include the supply of tangible resources such as water and sustainably produced crops and timber (provisioning services), and processes that regulate water and air quality, prevent natural hazards such as flooding and soil erosion, and mitigate climate change through storing and sequestering carbon (regulating services) (Dudley & Stolton, 2003; Brown et al, 2006; Campbell et al, 2008). Protected areas also provide cultural services, for example by supporting recreation and tourism, and maintaining cultural identity and a sense of place (Butcher Partners, 2005; Eagles & Hillel, 2008).

These services are underpinned by the role that sites play in supporting the preservation of basic ecological processes (e.g. nutrient cycling), fundamental in maintaining the overall functioning of natural systems (supporting services noted). Healthy and well-functioning ecosystems sustained within protected areas can increase not only the range of ecosystem services, but also the resilience of ecosystems to resist and adapt to disturbances (e.g. climate change) also beyond the site level (Stolton et al, 2008; Dudley et al, 2010).

The Natura 2000 network, while almost complete at the terrestrial level, has yet to be finalised for marine protected areas (MPAs), and much of the network is still not yet reaching favourable conservation status (FCS) (see Glossary in Annex I for definitions). The systematic assessment covering the reporting period from 2001 to 2006⁵ concluded that only 17 % of the assessments for both habitats and species were favourable though this is quite variable across the regions (see Figure on previous page).

More needs to be done to improve the ecological status of the network. A healthier Natura 2000 network will also lead to a higher level of benefits provision to society and the economy as well as be more resilient to environmental pressures including climate change.

⁴ IP/11/1376: Press Release: Environment: Major expansion of Europe's protected natural areas available via <http://europa.eu/rapid/>

⁵ COM(2009) 358 final. Composite Report on the Conservation Status of Habitat Types and Species as required under Article 17 of the Habitats Directive. Brussels. Member States report every six years on their progress in implementing the Directive and the status of habitats and species of Community interest.




II) Methodology for assessing EU wide benefits of Natura 2000

Methodological Framework

This study has employed an ecosystem services approach to assess the benefits delivered by the network, and to examine their value. By protecting Natura 2000 sites and requiring conservation action, the network should enhance the functioning of ecosystems, which in turn deliver benefits to society and the economy (Figure 3).


The Millennium Ecosystem Assessment (MA) provides a framework for categorising, assessing and valuing the services delivered by ecosystems. Sites deliver a range of provisioning, regulating and cultural services that enhance human welfare. These are underpinned by supporting services, which benefit people indirectly⁶.

Figure 3: Benefits of Natura 2000



Source: Adapted from Braat and ten Brink et al (2008)

⁶ Care needed to avoid double counting.

Figure 4: Contribution of Ecosystems and Biodiversity to Human Wellbeing


Source: TEEB 2011, adapted from Haines-Young and Potschin (2009) and Maltby (2009)

Some services are directly linked to species' detailed composition and diversity (e.g. pollination, cultural services). Others, like flood regulation, depend on the role of physical structures and processes at the ecosystem scale. These ecosystem services, in turn contribute to human wellbeing by providing a range of environmental, social and economic benefits – see Figure 4.

To examine the overall value of the multiple benefits delivered by Natura 2000 sites, a Total Economic Value framework (Figure 5) is used. This recognises that the values associated with the Natura 2000 sites result from their direct use by people (for example in the provision of food, fibre, fresh water and genetic resources, as well as cultural uses such as for recreation) as well as their indirect uses (for example in regulating air, water and climate). In addition, people derive non-use values from the existence of sites and their protection for future generations. It should be recognised that this framework captures only the value of Natura 2000 from an anthropocentric viewpoint – i.e. the benefits that sites provide to people – and that biodiversity has an intrinsic value that is independent of human thoughts and values. These intrinsic values – while an important motivation for establishing the network – cannot be captured by the ecosystem services framework and are not therefore estimated.

It should also be noted that, in general, ecosystem services assessments are still in a stage where their science base (ecology and economics) is still under development. The assessment carried out in this study built on the best science available to date (e.g., MA, TEEB) and relies conceptually on existing ES typologies and knowledge (e.g. on trade-offs, resilience etc.). As these concepts are still being refined, and the literature used were produced in different periods (and mostly before the MA and TEEB), a mismatch between data sources for the assessment (case study valuations) and conceptual (scientific) underpinning of the report may exist (e.g., double counting in some case studies, suboptimal research design, etc.). Furthermore, it should be acknowledged that this assessment is based on the current state of the world as understood by this evaluation. It is likely that the supply and demand drivers of Natura 2000 will change over time and therefore today's assessment may not be easily extrapolated to the future. These nuances should therefore be kept in mind, although the report strived to be on the frontline of the debates and data used by the most recent exercises, such as TEEB.

Figure 5: The Total Economic Value (TEV) framework in the context of Natura 2000



Source: White et al, 2011, adapted from Kettunen et al (2009), adapted from Pearce & Moran 1994

Benefit/Value transfer

This assignment has involved the development and application of benefits transfer methodologies (now increasingly termed ‘value transfer’), using existing valuation evidence of the benefits of Natura 2000 sites as a basis for estimating the benefits of the network as a whole. Benefit transfer involves the application of values obtained in one context (the ‘study site’) to estimate the value of benefits in another context (the ‘policy site’)⁷. It provides a cost-effective means of deriving overall value estimates, but needs to be applied with great caution, taking care to ensure that the values used are robust, relevant and applicable to the policy site.

Assessing Overall Benefits – Alternative Methods

The study used four different methods to assess the overall value of the benefits of Natura 2000 sites and to aggregate them to assess the overall benefits of the network: an ecosystem service-based, a territorial-based, a site-based and a habitat-based approach. The territorial-based approach proved too weak given data and methodology issues and not used in the final assessments. The table below provides a description of the three methods that were used in the final work, as well as their pros and cons.

⁷ There are different types of benefits transfer: unit benefit transfer – e.g. multiplying a mean unit value (per household or per hectare) from a similar site by the quantity of the good/ service at the site being assessed; adjusted unit benefit transfer; value function transfer and meta-analytic value function transfer. See the Full Report for further discussion and sources.



A range of **key methodological issues** have been taken into account in the course of assessment. They include the issue of gross and incremental benefits of Natura 2000, the additionality of benefits, opportunity costs and trade-offs, spatial variations in benefits and values, non linearity and thresholds, discounting, aggregation and scaling up, double-counting and ecological knock-on or ecological multiplier effects. These are discussed in detail in the Full Technical Report (<http://ec.europa.eu/environment/nature/natura2000/financing/>).

Table 1: Methods for Benefits Estimation and their Strengths and Weaknesses

Method	Description	Strengths	Weaknesses
Ecosystem Service Based	This approach focuses on the contribution of Natura 2000 to the delivery of individual ecosystem services, seeking to quantify and value each service.	Consistency of approach for valuing each individual service. By focusing on particular services, may provide relatively robust lower bound estimates of value of benefits.	Geographic variations in service delivery make estimation at network level difficult. Only certain services can be valued so likely to underestimate benefits of the network.
Site Based	Benefits estimates are available for a number of different Natura 2000 sites. These can be scaled up to estimate the benefits at network level.	Draws on data from a relatively large number of studies (though still small compared to optimal). Recognises and has the potential to account for the different characteristics of sites and the nature and value of services they deliver.	Difficulty of accounting for wide variations in estimates between sites (unless very large base data). Amalgamates estimates produced using different methods. Difficulty of knowing how available estimates relate to overall characteristics of network and providing a robust basis for upscaling.
Habitat Based	Site based estimates can be used to estimate per hectare values for individual habitats, which are then combined with data on extent of habitats at network level, to provide EU wide estimates.	Provides a logical basis for upscaling, as similar habitats are likely to deliver similar types of services across the network (although the value of many services varies significantly by location). Data are available on area of individual Natura 2000 habitats, providing a basis for upscaling.	Variations in service delivery can be expected within habitats, according to location. Difficulty of accounting for wide range of benefits estimates for certain habitats. Lack of estimates of benefits of some habitats. Amalgamates estimates produced using different methods.

Part B:

Deriving an aggregate total value of services from Natura 2000

III) The total Value of the Natura 2000 network – a first assessment

This section presents overall estimates of the value of the benefits delivered by the Natura 2000 network, based on aggregation of site-based and habitat-based data.

Site-based estimates of Natura 2000 benefits

Overview of approach

Various studies are available of the benefits provided by different Natura 2000 sites. These studies indicate that different sites deliver different benefits and that estimates of the value of these vary widely – this may reflect the value of the benefits themselves as well as the degree to which they can be valued comprehensively and accurately.

Compiling data at the site level provides a basis for scaling up across the network as a whole. Site based estimates can be pooled to give a range of per hectare values for sites. While different studies may focus on different services and benefits, reflecting the different characteristics and locations of sites and the services they deliver, this is not necessarily a problem if the individual studies are robust and provide a relatively complete and consistent approach to benefits estimation.

Advantages of this approach are that it enables a relatively large number of existing benefits estimates to be employed, and that it recognises the natural variations in sites and their characteristics and values. A key disadvantage is that it combines values from a range of different studies employing different methods and assumptions, whose consistency may therefore be questioned. Furthermore, scaling up from the site to the network level presents methodological issues and challenges, given the variability of site based estimates.

Available Benefits Estimates

An extensive review was undertaken of studies assessing the value of services delivered by Natura 2000 sites. The analysis focused on studies that:

1. **Cover a wider range of ecosystem services** provided by the sites in question, in order to enable a reasonably complete assessment of benefits. While data constraints often preclude comprehensive analysis of the value of ecosystem services, studies that focused on one or two services only were excluded from the assessment. Since most values identified covered certain services only, they are likely to provide a conservative estimate of the benefits of the network;
2. **Provide estimates of the annual per hectare value of benefits**, or enable such an estimate to be derived. Estimating benefits on a per hectare per annum basis provides a standardised basis for the analysis and upscaling of values.
3. **Relate to terrestrial and coastal sites only** – the benefits of marine sites are considered separately below.

The review provided 34 different estimates of the value of the benefits of Natura 2000 sites, from 20 different studies. A summary of studies and their value estimates is given in Annex II. This analysis revealed that the available estimates give a wide range of values for the benefits of Natura 2000 sites, ranging from just less than €50 per hectare per year to almost €20,000 per hectare per year. The range of values identified underscores that sites are not uniform, while estimates of the value of the services they deliver also vary according to the methods used and data available.

Variations in value estimates reflect differences in:

- The location and characteristics of different sites (including their condition, scarcity and substitutability);
- The ecosystem services delivered, which vary by habitat and location relative to people and natural resources;
- The value placed on those services by people and by markets;
- The extent to which studies have been able to estimate ecosystem service delivery and its value;
- The methods used in valuation, and the assumptions used in benefit estimation; and
- The role of non-use values which can form a significant share of the total value.

Although the available values have a wide geographical spreads, the majority come from North West Europe, particularly the UK and the Netherlands, which raises some concerns about their applicability to the network as a whole. While it is possible to take account of some variations between Member States when scaling up to the EU as a whole (e.g. by adjusting for variations in GDP), it is likely that the sample of values does not fully account for variations in ecosystem service delivery across the network.

Estimating the benefits for the EU27

Two methods are employed to upscale these estimates to the EU level:

- a. Simple upscaling based on mean and median per hectare values for sites;
- b. Upscaling of GDP adjusted mean and median per hectare values for sites.

Development of alternative approaches (including a typology of sites and the development of a benefit transfer function) was also explored, and is discussed below, but proved to be unworkable due to limitations in the data available.

The second approach was considered more robust and feasible for this study, and is therefore used. A discussion on other possible approaches is provided in the Full Technical Report (ten Brink et al, 2011).

Use of GDP adjusted per hectare values

The value of benefits can be expected to vary according to differences in income levels between Member States, which affect the value of ecosystem services and willingness to pay for them. Each of the site-based estimates was adjusted for differences in GDP per capita in Purchasing Power Standards, in order to provide income adjusted estimates of the value of benefits per hectare. The adjustment used Eurostat indices of national GDP per capita, on the basis that Natura 2000 sites provide benefits at the national level, and most studies estimate benefits to the national as well as the local population.



Because the available estimates are concentrated among higher income Member States, adjusting them for differences in GDP per capita reduces the overall benefit estimates (Table 2).

Table 2: Estimated benefits at EU27 based on up-scaling of GDP adjusted site based estimates

Basis for upscaling	Value per hectare (€)	Value EU27 (€M)
Mean	3,441	313,520
Median	2,447	222,951

Upscaling using these per hectare values gives overall benefit estimates of between **€223 billion and €314 billion** annually for the Natura 2000 network as a whole. This should be seen as a first illustrative estimate of the scale of the annual benefits and not as a robust precise result.



Habitat-based estimates of Natura 2000 benefits

Overview of approach

It is also possible to use estimates of the value of services delivered by different habitats as the basis for estimating the value of the benefits of the Natura 2000 network. Because similar habitats can be expected to deliver similar types of ecosystem services, we can expect the value of services to vary by habitat. Data are available for the area of different habitats in the network, and can be used as a basis for up-scaling habitat based values.

This is similar to the site-based method and involves compiling estimates of the value of benefits delivered by different habitats. These may be derived from studies focusing on a particular habitat (e.g. benefits of marine protected areas) or on particular sites dominated by a single habitat. A range of values can be derived for each habitat, and, combined with data for the area of each habitat covered by the network, used to provide estimates at the network scale.

This approach has some advantages in that there is likely to be some consistency in the types and levels of services delivered by a habitat, while good data on the areas of each habitat covered by the network are available. Disadvantages relate to the consistency and reliability of different benefits estimates, the likelihood that data will be unavailable for certain habitats, and the known variations in delivery of some services within habitats. For example, while some services such as climate regulation may be reasonably consistent between different forest sites, the value of others such as water purification will vary significantly according to the location of the forest (for example in relation to pollution sources, water supplies and centres of population). This presents challenges in extrapolating benefit estimates across the habitat as a whole.

Relevant data sources include: data from Cost of Policy Inaction (COPI) (Braat et al 2008; ten Brink et al., 2009 and the TEEB database (van der Ploeg et al 2010) and TEEB studies on value of services delivered by different habitats; and studies of individual sites as above where these have a predominant habitat or values broken down by habitat.

Available Benefits Estimates

The first step is to calculate the mean and median values for each habitat type identified through the literature review. To calculate the habitat values, the site based studies summarised above were grouped by broad habitat types, using the Habitat Directive Classification. This was not straightforward, as the studies reviewed did not use the Natura 2000 habitat classification system in their reports. Judgement was required to associate the habitat included in the studies reviewed with the Natura 2000 classification system. In addition, several of the values identified were based on studies of sites that contained more than one habitat type. When this was the case, the value contained in the study was assigned to the predominant habitat type for the site in question.

The mean and median value for Natura 2000 sites, by hectare, and adjusted for differences in GDP (from 2010 Eurostat figures), were calculated based on the entire range of values identified. The results of these calculations are presented in the table below.

Table 3: Natura 2000 habitat values, per hectare

GDP adjusted, 2011 €/Ha/year					
Habitat Directive Classification (Natura 2000 habitat code)	Count	Min	Max	Median	Mean
Coastal and Halophytic Habitats (1)	6	743	3,954	3,053	2,651
Coastal Sand Dunes and Inland Dunes (2)	2	3,863	9,849	6,856	6,856
Freshwater Habitats (3)	8	371	4,685	1,231	2,256
Temperate Heath and Scrub (4)	3	1,009	17,336	5,252	7,866
Sclerophyllous Scrub (Matorral) (5)	0	–	–	–	–
Natural and Semi-natural Grassland Formations (6)	5	77	5,875	1,156	1,898
Raised Bogs and Mires and Fens (7)	3	136	12,956	951	4,681
Rocky Habitats and Caves (8)	0	–	–	–	–
Forests (9)	5	347	4,969	924	2,309
All habitats	32	77	17,336	1,721	3,323

Source: TEEB 2011, adapted from Haines-Young and Potschin (2009) and Maltby (2009)

Estimating the benefits for the EU27

These per hectare values can be combined with data for the area of each habitat across the Natura 2000 network to estimate the value of benefits for the network as a whole.

Data on the area of each habitat in the Natura 2000 network was identified in Mùcher et al. (2009). This information is used to estimate the total value of the Natura 2000 network, by habitat, based on median, mean and mean excluding outlier values. The results of this analysis are presented in Table 4 below.

Estimates are made for 7 habitats for which values are available, and scaled up to the Natura 2000 network as a whole.

This method gives estimated values of between **€189 billion and €308 billion** per annum, depending on whether the median or mean values are used. The figures are slightly lower than for the site-based estimates, because the most widespread habitats (such as forests) have slightly lower estimated per hectare values than the average.

The outlook

The above first estimates offer order of magnitude value ranges for the gross benefits of the Natura 2000 network. These should be taken as illustrative estimates which can help communicate the economic value of the range of socio-economic co-benefits stemming from the ecosystems covered by the Natura 2000 network.

There is a clear need for further site based studies which are more geographically spread across the EU, that cover a wider range of ecosystem services and are done in a comparable manner which would help create an improved evidence base for future assessments – as well as being immediately useful to demonstrate benefits for the local to national to EU stakeholders. The details on the outlook is presented in part E.

Table 4: Estimated Natura 2000 habitat values

Habitat Directive Classification (Natura 2000 habitat code)	Estimated area (million ha)	Median	Mean
Coastal and Halophytic Habitats (1)	15.0	45,884	39,849
Coastal Sand Dunes and Inland Dunes (2)	1.5	9,993	9,993
Freshwater Habitats (3)	6.2	7,628	13,977
Temperate Heath and Scrub (4)	11.5	60,284	90,285
Sclerophyllous Scrub (Matorral) (5)	4.0	–	–
Natural and Semi-natural Grassland Formations (6)	11.6	13,373	21,964
Raised Bogs and Mires and Fens (7)	7.8	7,450	36,672
Rocky Habitats and Caves (8)	4.1	–	–
Forests (9)	29.4	27,189	67,956
Total (7 habitats)	83.0	171,802	280,695
Estimated Total for Natura 2000 Network (9 habitats)	91.1	188,587	308,118

Part C:

The value of different Ecosystem services from Natura 2000


IV) Overview of Ecosystem Services

The Natura 2000 network provides benefits to society and the economy through the delivery of different ecosystem services, with the importance of each service varying between sites, depending on site characteristics, location, and the type and level of interaction between the ecosystem and the social and economic systems or, to put it differently, between the ecological/green infrastructure (with Natura 2000 sites as core connected areas within a wider green infrastructure) and the economic and social structures. As noted above, the level of knowledge of the different ecosystem services varies and is changing fast as more attention is paid to the issue.

A study by Gantioler et al. (2010) explored what experts consider to be 'key' benefits associated with Natura 2000 (structured interviews were carried out with more than 110 individuals from 26 Member States, including representatives of national governments, NGOs, stakeholder groups and academia). The interviewees were requested to estimate (on a scale of 1 to 5) the importance of Natura 2000 in providing different ecosystem services – at local, national and global level – to obtain an indicative overview of the level of appreciation and the perceived relevance of Natura 2000 in providing those services. The results are presented in Figure 6 below.



Figure 6: Perceived relevance of Natura 2000 in providing different ecosystem services at local, national and global level (on a scale of 1-5)



Source: Gantioler et al, 2010

This was an exploratory assessment and is an interesting starting point for the discussions here. The values depend on the specific site and significant variations can be expected for each of the 'overall' judgements noted below.

The following sections summarises the assessment of the benefits of Natura 2000 network for a core set of ecosystem services selected for this study. This is a first attempt to develop an aggregate value for the Natura 2000 network as a whole. Given existing gaps in evidence and data, the site specific nature of many services, and a range of methodological challenges, these assessments of different services present different levels of answers. The aim here was to only present aggregate numbers for the network as a whole where sensible, and to note where approaches should only be seen as 'experimental'.

The value of carbon storage and sequestration – this is included first given the high policy relevance and since that the values assessed are relatively robust. Key insights on the benefits of natural hazard mitigation and climate adaptation, the value of tourism and recreation (this is a summary of a parallel study by Arcadis et al, 2011 forthcoming), water provision and purification, food-related provision (fish provision in marine areas and pollination and agricultural production in terrestrial areas), and health, identity and learning benefits.

V) Natura 2000's fundamental role in climate mitigation – The Carbon storage and sequestration benefits of Natura 2000

Introduction

In general carbon stock density appears to be relatively high across Europe (Campbell et al., 2008). Many Natura 2000 sites harbour several ecosystems that are important current stores of carbon and offer significant opportunities for further carbon sequestration, including sites located on forested lands, wetlands, agricultural lands, and marine and coastal ecosystems. In particular Northern European countries, where boreal forests are predominant, show much higher carbon storage potential in terms of high carbon density in the soil and biomass. Therefore, a careful assessment of carbon potential and economic consequences associated with Natura 2000 habitats may provide important insights on the cost-effective land-use policy and management practices on Natura 2000 sites, which in turn can influence ecosystem progress that affect greenhouse gas (GHG) fluxes⁸ over a period of several years to a few decades, and contribute to climate change mitigation and adaptation strategies in Europe.

Evidence and Results

A comprehensive economic valuation of carbon benefits provided by Natura 2000 sites needs a solid scientific base. The present estimation of the carbon benefits is built upon the 2003 IPCC Good Practice Guidance (GPG) *for Land Use, Land Use Change, and Forestry (LULUCF)*. The valuation framework was developed following three key steps:

Step 1. Characterisation of the status quo (SQ) or baseline scenario in 2010.

This step involves profiling the current carbon economic value provided by all Natura 2000 sites in Europe in a reference year (2010). To calculate the total carbon stocks by habitat type, the simplest and most practical way is to multiply carbon density (tC/ha) of each habitat type by the total area of the existing habitat. In our study, estimated carbon density stored by different habitat types was derived based on a review of the literature and selected from the studies that included habitat types most relevant to the Natura 2000 habitat classification. Furthermore, the carbon sequestration services are translated into monetary terms by applying a range of carbon prices to reflect the damages caused by different degrees of climate change impacts.

⁸ CO₂ fluxes between the atmosphere and ecosystems are primarily controlled by uptake through plant photosynthesis and release via respiration, decomposition and combustion of organic matter.

All in all, our valuation estimates indicate that the total carbon value of all Natura 2000 habitats as a whole⁹ lay between €607 and 1,130 billion in 2010, depending on the choice of carbon prices. These are values of the stock of carbon and not the annual sequestration rate. Among all others, the forest habitats contain the highest carbon value in the network, ranging between €318.3 and 610.1 billion in 2010. The second highest carbon value is contained in the dryland (grassland) system, ranging between €105.6 and 196.5 billion in 2010, followed by marine and inland water ecosystem, which account for €92 to 171 billion and €84.2 to 156.7 billion, respectively¹⁰.

Step 2. Characterisation of a future scenario by 2020 – the EU policy target year.

This step involves the study of policy-driven land use changes and the assessment of their respective impacts by 2020, in terms of changes in carbon stocks in the above ground biomass and below- ground soil organic matters. Given that CO₂, the most common GHG, is sequestered in biomass and soils in forests, wetlands and grasslands at higher rates than in cropping systems. A number of management practices on Natura 2000 sites that can result in an increase in soil organic carbon and carbon sequestered by biomass are identified.

Onsite measures that positively affect carbon fluxes include the restoration of wetlands, including peatlands, the improvement of grassland and the establishment of agro-forestry ecosystems. On the contrary, policies that passively manage the existing protected areas or encourage land conversions from grassland to croplands will cause the release of stocked CO₂ to the atmosphere and reduce carbon stored in the ecosystems.

⁹ Note that for the carbon analysis a low area of Natura 2000 network was used (51.5 million hectares, using 2009 data), given data availability and methodological needs. Data source: EEA data 2009 (<http://www.eea.europa.eu/data-and-maps/data/natura-2000-eunis-database>). This underlines that the results further below should be seen as a very conservative estimate.

¹⁰ In order to value the carbon sequestration services of Natura 2000 habitats in monetary terms, a range of carbon prices are applied to reflect the damages caused by different degrees of climate change impacts. In the present report, the European Commission values of €17.2/tonne in 2010 and €39/tonne in 2020 (EC, 2008 and DECC, 2009) are used as the lower values, and those building on a French study – €32 and €56/tonne in 2010 and 2020 respectively (Centre d'Analyse Stratégique, 2009) – as the higher values.

These considerations led us to focus on two types of possible future paths regarding Natura 2000 site management in Europe, i.e. (1) a **policy ON scenario**, where full Protected Area coverage (terrestrial PAs + fuller MPAs) with a move to full favourable conservation status will be evaluated; and (2) a **policy OFF scenario** where no additional action is taken and where some elements of degradation may occur across the Natura 2000 sites by 2020. More specifically, to assess the impacts of 'policy ON' scenario on carbon stocks, a separate evaluation is given for:

- the **quality improvement** of existing Natura 2000 sites, based on the net annual change of C-stock (tC/ha/yr) due to improved land-use management (IPCC, 2000). This is referred to as Option 1; and
- the **quantitative changes** of Natura 2000 site in terms of changing in land-use composition and conversions between different land uses, where the 'stock change method' (Penman et al. 2003) is used to estimate the economic consequences of a hypothetical EU policy which is aiming for **at least 10% increase in forest-protected area in all the Member States by 2020, with respect to their national forest coverage in 2010**. This is referred to as Option 2.

For both of these options the total carbon value provided by Natura 2000 sites in 2020 can be estimated by multiplying the estimated total carbon stocks in 2020 by the carbon price in that year.

On the contrary, if neither of the policy ON options were undertaken, this is then a Policy **OFF** – '**policy inaction**' scenario, where the economic gains from improved policies on Natura 2000 sites are not forthcoming. In particular, the policy OFF scenario refers to a scenario in which the EU will not provide any future investments in Natura 2000 habitat protection and management. As a consequence, certain degrees of natural degradation may occur on many sites and thus result in the release of CO₂ to the atmosphere or loss of carbon value. However, it is scientifically uncertain, whether and to what extent, Natura 2000 habitats may degrade in the context of policy inaction. For this reason, in the absence of reliable information, a zero rate of degradation is assumed, meaning that by 2020 the total quantity of carbon stocked in these habitats will remain the same as in 2010 (Status Quo).

Step 3. Interpretation of policy impacts and associated losses/gains on carbon value by comparing the selected policy scenarios and the SQ.

The results derived from both qualitative and quantitative evaluation of potential policy (ON and OFF scenarios) impacts can be integrated into cost-benefit analysis of the policy alternatives to provide important insights on cost-effectiveness of these policies. In Table 5 and Table 6, the estimated total carbon stocks and the respective economic values of Natura 2000 habitats, under different policy scenarios is summarised.

Table 5: Estimated total carbon stocks by Natura 2000 habitats (GtC)

Scenarios	Total	Marine Total	Inland Water Total	Dryland ESS Total	Cultivated ESs Total	Forest & Other Wood Land Total	Inland rocks, Scree, Sands, Permanent Snow & ice	Other land
Policy OFF Scenario in 2020	9.61	1.46	1.33	1.67	0.43	4.47	0.25	0.00
Scenario Policy ON-1 in 2020	9.78	1.46	1.33	1.74	0.45	4.55	0.25	0.00
Scenario Policy ON-2 in 2020	9.89	1.46	1.33	1.55	0.39	4.92	0.25	0.00

Note: see Table A4 in Annex 2 in FULL Technical Report for detailed results

**Table 6: Total Economic value of carbon services provided by Natura 2000 habitats
(Billion €, 2010)**

General habitats	Policy OFF – 2020		Policy ON_1: qualitative improvement – 2020		Policy ON_2: quantitative land-use changes – 2020	
	Lower bound	Upper bound	Lower bound	Upper bound	Lower bound	Upper bound
Marine Total	208.6	299.6	208.6	299.6	208.6	299.6
Inland Water Total	191.0	274.3	191.0	274.3	191.0	274.3
Dryland ESs Total	239.5	343.9	248.7	357.1	221.5	318.1
Cultivated ESS Total	62.2	89.3	64.5	92.6	55.6	79.8
Forest and Other Wood Land Total	639.7	918.6	651.8	936.0	703.7	1010.4
Inland rocks, Scree, Sands, Permanent Snow and ice	35.6	51.1	35.6	51.1	35.6	51.1
Other land	0.0	0.0	0.0	0.0	0.0	0.0
Total	1376.7	1976.8	1400.3	2010.6	1416.0	2033.3
Δ wrt Policy OFF (Δ %)	–	–	+23.6	+33.8	+39.3	+56.5

Note: see Table A6, A7 and A8 in Annex 2 in FULL Technical Report for detailed results

A policy scenario (Policy ON), where full Protected Area coverage (terrestrial PAs + fuller MPAs) with a move to full favourable conservation status is estimated to generate a gain of at least a total of 1.7-2.9 by 2020 compared to a policy inaction scenario (Policy OFF), where no additional action is taken to conserve the current Natura 2000 sites over the next decade.

VI) Natura 2000 as a tool for security: Natural hazards benefits and climate adaptation

Introduction

Among the wide range of benefits they provide, protected areas are known for their important role in mitigating the damaging impacts of natural disasters (e.g. TEEB, 2011; MA, 2005). In particular, protected areas are recognized to maintain healthy, intact and robust ecosystems, which help mitigate the impacts of disasters and restore destroyed or degraded areas (Mulongoy and Gidda, 2008). Protected areas play as well an important role in decreasing the vulnerability of communities to disasters and reducing their physical exposure to natural hazards, often providing them with livelihood resources to withstand and recover from crises (ibid).

In general, ecosystems affect both the probability and severity of extreme events, and they can moderate their effects. For example, inland waters, such as lakes and wetlands, are traditionally considered to be very important for the temporal regulation of water flow, mainly by accumulating water during wet periods (reducing peak flow). There is evidence that floodplain wetlands have the effect of reducing or delaying floods.

Natura 2000 sites can and have played a significant role in prevention and mitigation of extreme natural events. Moreover, due to population increase and climate change impacts, it is expected that the vulnerability of human settlements to natural hazards will increase in the future.

The potential of sites to control extreme events depends on the ecosystem types they host and their characteristics. For instance, an ecosystem's ability to mitigate avalanches is directly related to its forest cover and tree density; sites located along catchments areas (e.g. river slopes and floodplains) and coastal zones are likely to play a role in regulating water flows. Although the benefits arising from natural hazards risk reduction are very site-specific, well-functioning ecosystems in disaster-prone areas can offer efficient mitigation services, often at a lower cost than man-made measures.

Overall, growing attention is being paid to ecosystem-based solutions for natural hazards mitigation. Increasing evidence suggests that, in many cases, a degradation of natural ecosystems is likely to lead to exacerbated consequences of natural hazards (Dudley et al., 2010). Using ecosystem-based rather than man-made solutions has often proved to be significantly cost-efficient, and natural hazards protection measures are increasingly being incorporated into land-planning strategies. However, it has to be noted that the exact functioning of ecosystems in natural hazards mitigation is still insufficiently understood and needs to be improved (see e.g. TEEB, 2011; MA, 2005).



Evidence and results

Natural hazards have caused significant damage across the EU over time. Extreme events in Europe have led to over eighty thousand cases of premature mortality over the period 1980 to 2010. Around 15 million people in Europe have been affected over the period with an associated cost estimated at around €163 billion. This equates to an annual average damages of €7 billion/year.

Europe has suffered over 100 major damaging floods in recent years. It has been estimated that, since 1998, floods have resulted in about 700 fatalities, the displacement of about half a million people and at least €25 billion in insured economic losses (EEA, 2004).

It is also widely acknowledged that the flooding risk in Europe is increasing as a result of climate change – i.e. due to higher intensity of rainfall as well as rising sea levels (IPCC, 2001). Additionally, there has been a marked increase in the number of people and economic assets located in flood risk zones (European Commission, 2007). The value of the regulation that is provided by different ecosystems is therefore likely to be escalating, given an increase in human vulnerability to natural hazards (TEEB, 2010).

There are a number of studies which recognise the importance of Natura 2000 sites for natural hazards prevention. For instance, in the analysis of the Azoras Islands Natura 2000 site by Cruz and Benedicto (2009), the regulation of extreme events is ascribed the highest level of importance, although no explicit valuation exercise was possible. It is noted, however, that floods and landslides are very frequent in the area, and in 1997 caused 29 deaths and around €20 million in damages. Similarly, in Oaş-Gutâi Plateau and Igriş site in Romania and in Białowieża Forest in Poland flood protection has been assigned a high level of importance although, due to the lack of data, a valuation was not possible (Kazakova and Pop, 2009; Pabian and Jaroszewicz, 2009) – see also Box 1 below for some additional examples.

In general, the valuation of ecosystems' ability to mitigate natural disasters, such as floods, storms and avalanches, is a very complex issue. Due to the functional variability of the sites and other influencing factors, such as proximity and position of a site to human settlements, the actual delivery of natural hazards mitigation varies from area to area. Moreover, the amount of valuation studies and the degree of representative values that could be used for benefit transfer and wide approximation – especially for Natura 2000 sites – is very limited. As such, it has been difficult to identify examples in the EU where there is a direct linkage between protected areas and natural hazards control. Due to these difficulties, it was not possible, within the context of this study, to provide an estimate of the overall benefits of the Natura 2000 network related to natural hazards protection.

Box 1: Some examples of natural hazard mitigation services within the context of Natura 2000

In Kalkense Meersen Natura 2000 site, in Belgium, it has been estimated that the restoration of the original river landscape by means of wetlands and estuarine habitats restoration can bring flood mitigation benefits between €640,000–1,654,286 per annum (Arcadis Belgium et al., 2011 forthcoming).

With regard to flooding, the Natura 2000 network has an important role to play in particular in mountain areas, where floods often originate. Mountain areas are also generally more flood-prone due their topography, hence they are most likely to benefit directly from natural protection. Considering that many of Natura 2000 sites are located in mountain areas, the regulation of water discharges and of natural storage mechanism in these areas can benefit many river systems throughout Europe (EEA, 2010).

In the Alpine region in Switzerland the use of forests is recognised as a major component of disaster prevention. Today Swiss forests, are managed mainly for their protective function. (ISDR, 2004; Dudley et al., 2010)

VII) Natura 2000 as a motor of the economy/oil of society – The tourism and recreation benefits

Introduction

A related study 'Estimating the economic value of the benefits provided by the tourism/recreation and Employment supported by Natura 2000'¹¹ estimated the benefits of tourism, recreation and employment supported by the entire Natura 2000 Network. The economic value or the benefits provided by tourism and recreation (i.e. market on non-market benefits) that refer to use values, and the direct and indirect employment supported by the Natura 2000 network were taken into account.

In assessing the benefits of tourism and recreation, it is important to distinguish between:

- The **recreational benefits** derived by visitors to Natura 2000 sites – i.e. the value of the recreational experience itself; and
- The **economic impacts** that expenditures by these visitors deliver to local economies, by supporting employment and incomes. These are important benefits of the network, although they do not provide estimates of the value of recreational benefits themselves. It should be noted that tourism expenditures can provide important benefits for local economies, but, since most of this money would be spent elsewhere in the absence of Natura 2000, aggregate expenditures do not constitute an overall estimate of benefits at EU level.

Evidence and results

Estimates of the recreational benefits that Natura 2000 provides to visitors were made by transferring benefits from other studies, which have used travel cost and contingent valuation methods to estimate the consumer surplus per visit. The economic impacts from tourism and recreation were calculated based on multipliers that were generated by the consolidated input-output tables from Eurostat. Non-market benefits related to recreation, on the other hand, were calculated on the basis of a site-based approach. The overall employment opportunities provided by the Natura 2000 network were calculated based on a land-use approach and scaling-up on a per-hectare basis. The authors estimate that:

- the value of recreational visits to Natura 2000 sites is €5-9 billion per annum, based on estimates of visitors' willingness to pay;
- the total expenditure related to tourism and recreation supported by Natura 2000 was between around €50 and €85 billion in 2006;
- the expenditure exclusively related to the visitors who have affinity for Natura 2000 sites (i.e. around 21 % of visitors to Natura 2000) was between €9 and €20 billion in 2006, generated by around 350 million visitor days;
- The total expenditure provided by tourism and recreation supports between 4.5 and 8 million Full Time Employment (FTE) jobs. The benefits generated by the visitors with affinity to Natura 2000 would support from 800,000 to 2 million FTE jobs. This compares to a total of about 127 million FTE jobs in the EU27 (in 2009)¹², and about 13 million jobs in the tourism sector (in 2008)¹³.

¹¹ http://ec.europa.eu/environment/nature/natura2000/financing/docs/Estimating_economic_value.pdf

¹² Eurostat, Employment in Europe 2010 – Statistical annex: http://ec.europa.eu/employment_social/eie/statistical_annex_key_employment_indicators_en.html

¹³ Eurostat tourism database: http://epp.eurostat.ec.europa.eu/statistics_explained/index.php/Tourism_employment#Database





- Natura 2000 sites have supported on average about 12 million FTE jobs each year in the EU during the period 2006–2008. This includes about 1.5 million jobs in agriculture, 70,000 jobs in forestry, around 200,000 jobs in fishing, 3.1 million jobs in recreation (excluding employment generated by hotels and restaurants), and 7 million jobs in other industries.

According to Eurostat, median gross annual earnings of full-time employees across all industries in the EU were € 12,236 in 2006. Taking this number as a basis, 11,870,000 jobs supported by the Natura 2000 network provide incomes of about € 145 billion per year. This must be considered as a rough estimate for two main reasons:

- The overall employment supported by the Natura 2000 network was estimated by scaling up data related to the dominant activities performed in a site, and stated in the Natura 2000 database. The authors state that '[the] estimates are subject to a relatively high degree of uncertainty, given the relatively small information basis from which the estimates were drawn and the multiple uncertainties related to the data gathering process.'
- The earnings figures applied (€ 12,236 per FTE) do not take account of the spatial distribution of Natura 2000 sites in the EU. In order to refine the results, one would have to calculate the economic benefits provided by direct and indirect jobs on a Member State basis, taking account of income differences in relation to the number of people employed (jobs supported).

VIII) Natura 2000 and ‘free’ resources for/value for money in the economy and society: Water purification and supply benefits (and waste)

Introduction

Water purification and provision are important ecosystem services that are provided by natural ecosystems, including protected areas such as Natura 2000. The economic value of water purification and provision will vary in each case depending on the type of ecosystem: in general, ecosystems that have intact groundcover and root systems are highly effective in improving water quality (Brauman et al 2007).

While it has not been methodologically feasible, given the site specificity of the benefits, to develop a robust EU wide assessment of the benefits of the Natura network for water purification and provision, it is clear from case examples that the Natura 2000 network can lead to cost-effective means of water purification and supply, offering significant savings over man-made substitutes.

Evidence and results

A number of major European cities, including Munich, Berlin, Vienna, Oslo, Madrid, Sofia, Rome, and Barcelona all benefit from natural filtration in different ways. These municipalities save money on water treatment due to natural treatment from the ecosystems. The savings can be passed on to consumers, resulting in lower utility costs for EU residents – see table below.

Information from the four European cities of Berlin, Vienna, Oslo and Munich allows an illustration of the benefits of protected areas for water purification and provision. Using benefit transfer, it can be estimated that the annual economic benefits of water purification are between €7 and €16 million and of water provision between €12 and €91 million per city. The average per capita benefits are between €15 and €45 per year for both water purification and provision combined in the four European cities analysed. This compares to average household water bills of €200 per year in the case of Germany.

This underlines that benefits can be indeed significant, and lead to substantial actual and potential cost savings from ecosystem based water purification and provision, both for companies (in terms of reduced operational costs) and citizens (reduced water bills). It will be important for cities to explore the role of natural capital (protected areas, wider green infrastructure) in the purification and provision of water and ensure that such considerations are integrated in the water management plans required under the Water Framework Directive.

Other examples of water purification and provisioning benefits, within and outside Europe, are noted in the table below. The variation between these values is accounted for by the different ecological functions of the ecosystems, the varying interactions with economic and social systems, and how the supply and demand for the services relate.



Table 7: Economic value of water filtration benefits from protected areas in Munich, Vienna, Berlin and Oslo.

City	Method of protection	Total area protected (hectares)	Land use	Amount of water supplied	Approximate number of people served	Benefits	Estimated annual value of water filtration based on m ³ produced	Estimated annual value of water provision based on m ³ produced
Munich	Protected areas and conversion to organic agriculture	6,000	1/3 agriculture, 2/3 forest	301,000 m ³ per day	1 million (80% of the city)	Decreased pesticide and chemical residues No treatment required	€ 8,624,915	€ 12,635,211 – € 47,168,232
Vienna	Strict protection, Vienna Water Charter	Over 60,000	All protected forest	400,000 m ³ per day	1.7 million (entire city)	No water treatment required	€ 11,461,681	€ 16,790,978 – € 62,721,903
Berlin	Groundwater protection zones	23,000 (1/3 of the city of Berlin)	Urban landscape, 40% 'green areas'	585,000 m ³ per day	3.5 million (entire city)	Less contamination	€ 16,762,709	€ 24,556,805 – € 91,730,783
Oslo	Landscape protection area	25,200	All protected forest and lakes	250,000 m ³ per day	455,000 (85% of the city)	Minimal treatment required	€ 7,163,551	€ 10,494,361 – € 39,201,189

Table 8: Overview of valuation studies – water purification and provision

Source	Original Value	2009 Value (EUR) ¹	Valuation Method	Location	Biome
Water purification					
Brenner Guillermo, J. 2007	403 USD/ha/yr	527	Benefit Transfer	Spain	Forests
Brenner Guillermo, J. 2007	3,191 USD/ha/yr	4,173	Benefit Transfer	Spain	Freshwater
Cruz, A. de la and J. Benedicto 2009	18,1 EUR/ha/yr	27	Replacement Cost	Portugal	Temperate forest and grassland
Water provision					
Cruz, A. de la and J. Benedicto 2009	99.7 EUR/ha/yr	122.3	Replacement Cost	Portugal	Temperate forest and grassland
Butcher Partners Limited 2006	39.8 NZD/ha/yr	22.9	Avoided Cost	New Zealand	Grasslands
Anielski, M. and S.J. Wilson 2005	0.076 CAD/ha/yr	0.06	Direct market pricing	Canada	Temperate and boreal forests

¹Adjusted by purchasing power parities and inflation

Sources: TEEB database, van der Ploeg and de Groot, 2010

Significantly more empirical research is needed to estimate the economic benefits of water-related ecosystem services provided by the Natura 2000 network to any level of robustness. The Full Technical Report can be consulted for an experimental assessment, which was carried out within this study¹⁴.

Work currently being undertaken by the European Commission's Joint Research Centre (JRC) on an EU-wide ecosystem service assessment (Atlas of Ecosystem Services, see Maes et al. 2011) could provide a means to link spatial data, such as the availability of water-related ecosystem services, to socio-economic data, and thus allow to account for demand-side characteristics at the local scale. In this way, a fine-tuned assessment of the water-related economic benefits of the Natura 2000 network seems possible. In particular, research is needed in two major fields:

- Primary valuation of (water-related) ecosystem services in protected area contexts. To date, the dependence of people (or water utilities) on protected areas can often only be estimated by analysing the design of relevant PES schemes. More primary research is needed on the dependence of communities on hydrological systems in protected areas. The use of Geographical Information Systems (GIS) can be helpful in this context.
- The EEA Land Ecosystem Accounts (LEAC) can provide a means to locate hydrological systems which are of high value to people. The work done by JRC is a first step in the identification of aquatic ecosystem services on a large geographical scale. Future work will need to include demand-side characteristics to take account of value differences in water scarce or water abundant regions, respectively.

¹⁴ Based on a very simplified extrapolation (exploratory assessment), the estimated annual value of natural water purification provided by forest and freshwater habitats in the Natura 2000 network could be estimated at €2.2–€25 billion and the estimated annual value of freshwater provided by the entire Natura 2000 network could be in the order of €2.8–€3.2 billion. These ranges should be seen as an experimental assessment and not formally used.

IX) Natura 2000 and food: Marine protected areas and fish, and terrestrial protected areas, pollination and agriculture

Food security and provision: Marine Protected Areas

Introduction – Marine Protected Areas

The Natura 2000 network is still developing in the marine environment, and has faced practical and conceptual challenges such as a lack of data on seabed habitats and identifying representative areas for mobile species. Nevertheless Natura 2000 designations are in place in coastal, inshore and offshore areas, and some of these have been subject to different types of economic analysis.

Current work means that an assessment of the habitat areas covered by a complete marine Natura 2000 network may be possible in 5 years.¹⁵ At the same time, increasing effort in marine valuation is creating an evidence base which can be used, along with appropriate assumptions and judgement, to assess the values attributable to the marine Natura 2000 network. Nevertheless, the lack of monetary evidence for many impacts, and even the lack of non-monetary quantitative evidence, remains a major challenge, in particular for individual sites. One approach to the lack of evidence on impacts has been to use expert judgement to plug data gaps (as in the Impact Assessment for the UK Marine Bill); other

impact assessments have focused on quantifiable costs and limited consideration of benefits primarily to qualitative descriptions (as in the individual Impact Assessments for specific UK MPAs).

Extrapolating from the UK results to the EU level can only be very approximate, because the figures are based on value estimates for UK seas, and because of a lack of information about the specific network and its habitats. The UK results show that the final estimate is quite sensitive to the details of network configuration: the values estimated ranged from about €71/ha per year to €132/ha year depending on the designation strategy.

To be conservative, extrapolation was based on the lowest value network, which provides general protection with somewhat increased representation of OSPAR habitats. In terms of annual equivalents, the values are approximately €1.4–1.5 billion per year for the current area of protection (4.7%), €3.0–3.2 billion per year for protection of 10% of sea area, and €6.0–6.5 billion per year for protection of 20%. The higher figures apply to stronger protection measures. They are only approximate annual equivalents and in fact the initial annual values would be lower, rising to higher values as the protection reaches its full impact on habitats and services. An EU network with stronger focus on particularly valuable habitats would be expected to give higher values.

Fisheries

The influence of marine protected areas on fisheries is a controversial topic. Natura 2000 management measures are likely to lead to a reduction and/or change (but in most cases not elimination) of fishing pressures. This may result in initial decline in catches from the site itself, but that could

¹⁵ Doug Evans, European Topic Centre for Biodiversity, pers comm. 20/7/11



enhance local populations and recruitment processes, and potentially even improve carrying capacities through effects on habitats. Fish can move out of the site to sustain or increase yields of nearby fisheries. So the Marine Protected Areas created through the network may have positive effects on overexploited fish stocks generally. Closed areas can already be used in fisheries management as a means of allowing overexploited stocks to recover and enhancing fishery productivity. On the other hand, if the fishing effort reduction within the site is simply displaced to increase efforts in adjoining areas, this could have negative impacts. The extent of these impacts is extremely difficult to predict, because of uncertainty about four key factors:

- The extent and location of the network;
- The level of protection, and in particular the types of fishing that will/will not be allowed in certain areas, and the efficacy of enforcement;
- The ecological relationships governing the resulting impact on fisheries, including the importance of reserve sizes and network effects; and
- The effectiveness of the revised Common Fisheries Policy (CFP) in controlling any displacement of fishing effort from protected areas, and more generally returning stocks to Maximum Sustainable Yield (MSY) levels.

Longer term, these complexities are compounded by potential fish-species range shifts in response to climate change.

Although fisheries productivity can be valued at regional or national levels, identifying the contribution of specific sites is difficult. Existing evidence is patchy and it is not possible to draw firm conclusions about the marginal impact of protecting sites. The fisheries benefits (or costs) of marine reserves will depend on management outside the reserves. Generally, benefits arise in particular where there is high effort prior to reserve implementation: if there are effective effort control mechanisms in place, fisheries' benefits from reserves may be small.

This complicates assessment, not least because fisheries management is dynamic: in particular, it is difficult to estimate how successful current attempts to reform the EU CFP, and allow European fisheries to recover from decades of overexploitation, may be. Beare et al (2010) documents the change to fish stocks in the North Sea as a result of the effective suspension of commercial fisheries during World War II. A dramatic change in age composition is observed. Recruitment to fish populations does not respond as dramatically as age structure, which is likely because other environmental conditions also influence it, and the effective closed period was not long enough for fish age structures to take effect on reproduction. The paper concludes that, had fishing been prohibited for a longer period of time than the six years of the Second World War, a population equilibrium with a higher proportion of older fish would have been


established. Maintaining such an equilibrium would have likely allowed a higher sustainable yield value, even if the total biomass catch was the same.

However, the management-dependent nature of fisheries benefits can be considered in a very simplified form by basic bio-economic modelling of fishery production, for example as assessed at the European level in the context of avoiding illegal, unreported and unregulated (IUU) fishing by Eftec (2008). The research models the influence on fisheries of IUU fishing, through dynamic bio-economic models specified across Large Marine Ecosystems (LMEs; e.g. North Sea, Celtic-Biscay Shelf), for commercial groups of fish species (e.g. Tuna and Billfishes, Cod-likes). This specification for LMEs and commercial groups avoids some (but not all) of the problems associated with competition among stocks and questions of achieving maximum sustainable yield (MSY) for individual stocks simultaneously.

It is possible to adapt the above method to assess the impacts of reducing effort, allowing stocks to recover. If it is assumed that protection of the Natura 2000 network can be represented by a 10% reduction in fishing effort – i.e. that fishing effort falls in the protected areas and is not simply displaced outside – then the models predict the results presented in the figure below. Catches at first fall (due to lower effort) but rapidly increase (due to increased stock sizes). Not all fish stocks are modelled – those included represent 46% of EU landings. If the non-modelled stocks respond in similar fashion, it might be expected roughly double the value, i.e. a total of approximately €1 billion per year after 20 years.

These estimates can be criticised on a number of grounds. They assume that the only source of reduction in fishing effort arises through Natura 2000 protection, and this is unrealistic given the on-going reform of the CFP. Further, they do not address the possible impacts of changes in carrying capacities or improvements in age structure. Off-site export of fish biomass is considered, but only approximately, in that the models effectively assume perfectly mixed stocks (the models are not spatial). Possible price changes are ignored. At best, therefore, these results might be viewed as indicative of the order of magnitude of potential for fisheries benefits to be achieved through Natura 2000 designations. To derive better estimates, it would be necessary to consider spatial models with more detailed representation of fish stocks and reproduction, as well as the spatial distribution of fishing effort, in conjunction with consideration of the reformed CFP. This would be a major undertaking, well beyond the scope of the present work. Perhaps the best approach would be to use Ecopath With Ecosim models (www.ecopath.org) for the marine systems.

Figure 7: Possible change in annual fishing values arising from reduced fishing effort associated with Natura 2000 protection



Food security and provision: Terrestrial Protected Areas

Pollination

Pollination represents an essential ecosystem service for human wellbeing, being a key ecological process on which natural and agricultural systems depend (e.g. TEEB, 2011; Millenium Ecosystem Assessment, 2005; Balmford et al, 2008). It is estimated that insect pollinators are directly responsible for 9.5% (around €153 billion) of the total value of the world's agricultural food production in 2005 (Gallai et al. 2009). Insect pollination is also estimated to increase the yields of 75 globally important crops and is responsible for an estimated 35% of world crop production (Klein et al., 2007).

Protected areas provide habitats and breeding grounds for pollinating insects and other species with economic and/or subsistence value (TEEB, 2011). The available area of natural habitat has a significant influence on pollinator species richness, abundance, and pollinator community composition. Habitat area in the neighbourhood of crop fields has been

found to be strongly related to a direct measure of the pollination service measured here in terms of pollen deposition provided by bees. Hence, as a network of natural and semi-natural habitats, Natura 2000 has a significant role to play in securing continuous provision of pollinating service in the EU.

The importance of Natura 2000 in providing pollination services has also been recognised by key stakeholders. In a survey assessment carried out to estimate the level of appreciation and awareness of Natura 2000 related ecosystem services, pollination was identified as one of the most relevant ecosystem services (Gantioier et al., 2010). However, from the existing evidence on pollination it is very difficult to provide any quantitative or monetary value of the benefits stemming from the Natura 2000 network. This is due to the fact that there is generally very sparse evidence on the values of pollination, especially in the context of Europe or protected areas, and where there is it tends not to differentiate between pollination from protected areas or from wider green infrastructure. Box 2 below presents selected examples of valuation studies in the EU, and Box 3 illustrates the potential of mapping this ecosystem service.


Box 2: Pollination values

Examples from the EU context

- The annual economic value of insect-pollinated crops in the EU-25 is about €14,2 billion (approximately 10% of the annual economic value for all food production in 2005). The number for global agricultural production amounts to €153 billion. (Gallai et al., 2009; for a more detailed analysis of this study see Annex 3)
- Using the methods of Gallai et al. (2009), the United Kingdom's National Ecosystem Assessment estimated the economic value of biotic pollination as a contribution to crop market value in 2007 at €629 million (England: €532 million, Northern Ireland: €28 million, Scotland: €69 million, Wales: unknown) (UK NEA, 2011)
- A recent EEA report (EEA, 2010) identifies the importance of natural pollination, particularly for alpine herbs, forests and semi-natural grasslands. Although the actual importance of pollination in the mountain ecosystems remains poorly known, it is important to acknowledge this in the context of our study – considering that many Natura 2000 sites are located in the mountain areas.
- Klein et al. (2007) found that the production of 87 out of 115 leading global crops (representing up to 35% of the global food supply) were increased by animal pollination.

Significant work in the area of ecosystem services mapping has been done by the European Joint Research Centre. In its recent Scientific and Technical Report (JRC, 2011) alongside maps of other ecosystem services, a map based on an indicator for pollination potential has been developed. In this regard, dependency ratios from Klein et al. (2007), visitation rates of pollinators based on distance relationships from Ricketts et al. (2008) and a spatial distribution of crops from Grizzetti et al. (2007) were used. From this data maps of 'the pollination potential or the capacity of natural ecosystem to provide pollination services to croplands' were constructed at the aggregated level and at more detailed 1 km resolution (see figure below). Ideally, future mapping exercises could combine ecosystem services mapping with Natura 2000 maps to better identify the services provided by the Natura 2000 sites.

Figure 7: Pollination potential, 1km resolution



Source: JRC, 2011

Agriculture and Natura 2000

Agro-ecosystems directly contribute to the provision of food for human consumption through supporting the world's agriculture. Biodiversity and ecosystems have also an indirect role in the world's food supply by, inter alia, allowing nutrient and water cycling or soil formation – see Table below, outlining the biodiversity benefits to agriculture.

Protected areas, such as the Natura 2000 sites, are often managed under agricultural schemes while still contributing to the principle of sustainable development and nature conservation. Contrary to the widely-held misunderstanding, designation of Natura 2000 sites does not aim to put all human activity on hold. In fact, many of the Natura 2000 sites are valuable because of the way they have been managed before their designation, and it is often desirable to continue with these activities to maintain the area's species and habitats in favourable conservation status. As such, there is a great interest in finding solutions that ensure Natura 2000 farmland remains productive, while at the same time maintaining and ideally improving the natural environment.

From this perspective, organic agriculture represents a promising agricultural management option for some Natura 2000 sites and protected areas under agricultural land-use. Although it does not necessarily imply high nature conservation value, it can offer clear benefits for biodiversity when compared to conventional forms of agriculture. Due to lower cultivation intensities and a bigger share of natural areas, more indigenous species are present in sites under organic farming, which in turn creates more intact and better-functioning ecosystems.

It is also likely that support for Natura 2000 and High Nature Value (HNV) farming in the EU offers significant synergies. However it is currently difficult to determine their potential overlap. It is known that the Natura 2000 network is protecting a significant portion of HNV farming area, especially parts that are of recognised biodiversity quality (for further information see Paracchini, 2008). Conversely, HNV farming directly benefits the conservation of Natura 2000 farmland habitats, being either within actual sites or in the wider countryside.

From the current available data it is difficult to estimate the portion of agricultural output directly attributable to the Natura 2000 network. As of now, it is possible to estimate only the portion of Natura 2000 area under agricultural use. Integrating the Natura 2000 spatial data with the Farm Accountancy Data Network (FADN) would allow a better determination of the agricultural output derived from Natura 2000 sites. Such integration is a pre-condition for any future estimation of agricultural benefits related to the Network.

It has to be noted that Natura 2000 farming also plays a significant role in the maintenance of local breeds and local plant and tree varieties adapted to valuable semi-natural habitats, as they are likely to play an essential role in future agricultural adaptation strategies. Replacement of these species by a smaller number of more productive breeds and crop varieties, which has been underway in the last decades, has proved to be one of the causes of biodiversity degradation and led to an increased vulnerability of the sector to external pressures such as climate change.

Table 9: Biodiversity benefits to agriculture through ecosystem services

Provisioning	Regulating	Supporting	Cultural
• Food and nutrients • Fuel • Animal feed • Medicines • Fibres and cloth • Materials for industry • Genetic material for improved varieties and yields • Pest resistance	• Pest regulation • Erosion control • Climate regulation • Natural Hazard regulation (droughts, floods and fire) • Pollination	• Soil formation • Soil protection • Nutrient cycling • Water cycling	• Sacred groves as food and water sources • Agricultural lifestyle varieties • Genetic material reservoirs • Pollinator sanctuaries

Adapted from UNEP, 2007. *Global Environment Outlook. GEO 4: environment for development. United Nations Environment Programme. Malta. p.172.*

Source: Adapter from UNEP, 2007

See Box below for some examples of the positive interrelation between agriculture and Natura 2000.

Box 4: Some Examples of Farming in the Context of Natura 2000

Being the first major farming for conservation project in Ireland, the Burren LIFE Project seems to offer a good 'value-for-money' solution with minimum estimated economic return of 235 %. (Rensburg et al, 2009)

Organic agriculture has been recognised as a particularly useful option within the Mount Etna national park in Sicily and the Sneznik regional park in Slovenia (EEA, 2010)

Traditional agriculture, and primarily sheep farming, has significantly contributed to the well-preserved and stable conservation status of habitats, flora and vegetation in the Island of Pag, in Croatia. Agriculture and conservation here co-exist, facilitating the production of the traditional cheese of Pag, and hence contributing to the local economy. The continuity of this situation is in the interest of the local population. (Sundseth, undated)

In a Rhön grassland area in south east Germany, mostly included in Natura 2000, an infrastructure for locally produced sheep products has been developed. Mowing and grazing through the use of sheep helped with site management, while a market for locally produced Natura 2000 products has been established. (Sundseth, undated)

X) Natural 2000 and our health, identity and learning

The role of ecosystems in supporting human health is manifold. First, naturally functioning ecosystems can regulate the range and number of species that are dangerous to human health. For example, a number of species (e.g. insects) are known to be vectors of human diseases (e.g. malaria, dengue fever, Lyme disease etc.). Disease control is usually more crucial in some developing countries. Nevertheless, also in the EU the well functioning of ecosystems can be beneficial in keeping the populations of some species under control, e.g. affecting competition on resources and predation.

Natural ecosystems are also a source of potential discoveries for pharmaceutical firms, although the benefits in the European context are rather limited in comparison to developing countries. For example, Gantioler et al (2010), from a review of existing evidence and interviews with national stakeholders, found that the role of Natura 2000 in preserving genetic and species diversity was recognised to be of high importance, but that the value of actual benefits gained from using sites for food, fibre, medicines and pharmaceuticals is currently low. Indeed, while Natura 2000 offers some potential for new commercial discoveries, there is no evidence of current interest in Natura 2000 sites as a resource for bio-prospecting.





Furthermore, natural ecosystems are known to play an important role in supporting physical and mental health by providing possibilities for outdoors activities, recreation and relaxation. Protecting the diversity of species and habitats helps to maintain a wider variety of possibilities for recreation, e.g. providing different natural settings and more opportunities for wildlife watching.

Finally, it is to be noted that ecosystems also play a positive role in protecting human health via a number of other functions, e.g. through the mitigation of natural hazards, and particularly by maintaining air quality. Ecosystems help to regulate air quality by removing contaminants, through physical processes such as filtration and biological processes such as decomposition and assimilation. Natural vegetation, especially trees and woodlands, improves air quality through the uptake, transport and assimilation of a wide range of gaseous and particulate air pollutants. Air quality regulation is especially supported by the maintenance and management of healthy forests with diverse vegetation structures and features increasing the surface area for the removal of pollutants.

Furthermore, access to natural compounds also plays a significant role in modern pharmaceutical research and development. It has been estimated that 25 of the drugs sold in developed countries and 75 of those sold in developing countries were developed using natural compounds (Pearce and Puroshothamon, 1992), demonstrating that biodiversity is of value to pharmaceutical firms in their efforts to develop new drugs.

Part D:

Realising the Benefits of Natura 2000

XI) Realising the Benefits: restoration and conservation for biodiversity and co-benefits.

A Completed and well-managed Natura 2000 network delivers the most co-benefits

As outlined in Part A, the Natura 2000 network is to a large extent complete for terrestrial areas with the focus now moving increasingly towards management and restoration activities (see below). On the contrary, the marine component of the network still requires more attention. It is commonly acknowledged that, despite increasing threats to the marine environment, progress in establishing marine protected areas (MPAs) has been very slow, particularly for the high seas. Globally it has been estimated that only 0.7 % of world's oceans are currently designated as MPAs (World Database

on Marine Protected Areas 2011). In the EU, marine areas currently account for close to 20% of the total Natura 2000 network with significant gaps still existing, especially in off-shore waters (Natura 2000 Barometer 2010).

Increasing the coverage of MPAs, including completing the Natura 2000 network in marine areas, is important also from the perspective of delivering socio-economic benefits. Research shows that appropriately designed and managed MPAs can provide a good conservation and sustainable management strategy for a range of species, particularly fish (Kettunen et al 2011). It has been estimated that conserving 20-30% of global oceans through MPAs could create a million jobs, sustain fish catch worth US\$70–80 billion/year and ecosystem services with a gross value of roughly US\$4.5–6.7 trillion/year (Balmford et al 2004). Furthermore, the improved ecosystems' health and conservation status of both terrestrial and marine protected areas and of the wider



Natura 2000 network is understood to improve the resilience of the functioning of the ecosystems – i.e. their ability to withstand pressures (e.g. climate change, pollution). This is expected to improve service provision (with improved health/connectivity) or reduce the loss of service provision, in light of climate change or other pressures risking degrading the ecosystem health.

Designation of areas valuable for biodiversity as Natura 2000 sites does not automatically guarantee that their favourable conservation status is maintained or restored. While designation is a valuable first step, appropriate management and restoration of Natura 2000 sites is needed to ensure that the conservation objectives – and related socio-economic co-benefits – are reached in practice. This requires sufficient financial resources, capacity to carry out management activities in an effective manner and continued support from both stakeholders and decision-makers alike. This report clearly illustrates that a well-managed Natura 2000 network can provide multiple benefits to both biodiversity and people. Without appropriate management regime, however, the effectiveness of Natura 2000 network is significantly reduced, undermining the supply of ecosystem services as well as conservation.

In parallel to this study assessing the total economic value of the benefits provided by the whole Natura 2000 network, work has taken place to identify the total economic value of the changes to ecosystem services as a result of taking conservation measures in Natura 2000 sites (Arcadis Belgium et al., 2011). The study has developed a tool specifically to value the changes in ecosystem services resulting from conservation measures taken in designated Natura 2000 sites. The tool has been tested with 11 sites reflecting a range of different geographical areas, habitat types and socio-economic circumstances across the EU and candidate countries.





For the majority of the sites the tool has produced results with a moderate level of confidence that shows that conservation measures deliver changes to ecosystem services that benefit people. As well as enhancing genetic and species diversity, in general, the conservation measures studied:

- Enhance cultural ecosystem services, including non-use value for landscapes and biodiversity, and visitor values;
- Sometimes increase carbon storage, although evidence that could be applied was limited to inter-tidal and forest habitats;
- Are expected to have positive impacts on regulating services. This area often lacks specific evidence (e.g. on air quality or erosion control), but some water quality and quantity regulation benefits could be valued;
- Can sometimes reduce provisioning services (e.g. reducing agricultural intensity) and sometimes increase them (e.g. maintaining or introducing grazing to maintain specific habitats).

The study also highlighted opportunity costs related to conservation and the implementation of management measures. In several cases reductions in intensity of provisioning services (e.g. agricultural outputs, timber) due to implementing a conservation regime were noted. However, when wider ecosystem services evidence (e.g. regulating services like carbon storage, and cultural services like landscape value) were taken into account, measures for which analysis was complete showed, as far as could have been assessed, net positive values for undertaking conservation measures.

Engaging stakeholders: attracting more support and resources

The ultimate success of the Natura 2000 network in reaching its set goals depends on the engagement of all relevant stakeholders in maintaining the network. Demonstrating socio-economic benefits arising from the management of the site can help to gain support among the different stakeholders. Furthermore, the assessment of these benefits can help to identify and address possible conflicts in a more precise and fair manner.

Demonstrating benefits can pave the way towards incentivising and creating new partnerships for conservation and/or sustainable use of Natura 2000 sites. In particular, it can help to provide new and innovative funding sources for site management (Kettunen et al. 2011, Gantioler et al. 2011).

It has in fact been estimated that the management of Natura 2000 would require around €5.8 billion/year (Gantioler et al., 2010). It is difficult to determine the overall funding available to support the network. However, the financial allocations for Natura 2000 from the EU budget are between €550–1,150 million/year (Kettunen et al. 2011). The estimated figures

represent only 9–19% of the financing needs. Existing financial resources are not yet adequate to ensure effective management and restoration of the network. This conclusion is also supported by the recent assessment of the status of the network, which concluded that only 17% of assessments of both the habitats and species conserved under the Habitats Directive were favourable (see Chapter 2). Consequently, it appears that the lack of adequate resources currently poses a significant risk to reaching the conservation objectives and also diminishes the amount of associated co-benefits from Natura 2000 sites.

In light of the current economic crisis, there is a clear risk that investment in protected areas may lose momentum and its places within the EU political agenda, and hence that fewer resources are made available for their management and protection. Whereas public funding will need to continue to be the main source of investment it will also be important to identify new and innovative sources of funding to ensure effective functioning of the network. These could include, for example:

Broader access to existing public funds (e.g. EU funding): Existing public funding for biodiversity conservation is known to be very scarce. However, significantly more resources are



available to support environmentally sustainable development of different regions and localities. Identifying and demonstrating links between the Natura 2000 network (e.g. its individual sites) and broader socio-economic benefits, such as natural risk management, climate change mitigation and adaptation, water management and tourism, can facilitate the access to a broader set of EU funds, including funds dedicated to support regional and rural development (i.e. European Regional Development Fund, European Social Fund, European Fisheries Fund and European Fund for Agriculture and Rural Development).

Payments for environmental services (PES): Evaluating the benefits of conservation can attract funds from stakeholders benefiting from the ecosystem services provided by Natura 2000 areas. The benefits of habitat conservation to water retention and quality in an area can form a basis for establishing a payment scheme where the users of water contribute to maintaining, managing and/or restoring the Natura 2000 site and its natural abilities to regulate the water flow. Similarly, as demonstrated in the study, Natura 2000 areas play a significant role in both storing and capturing carbon, to help mitigate climate change. Depending on future developments, this might make (certain) Natura 2000 areas eligible to receive funding from carbon offsetting schemes.

Revenues from certified products and/or sustainable tourism: With the markets for sustainably produced and/or biodiversity-friendly products increasing, there are also increasing opportunities to develop markets for certified products from or associated with Natura 2000 areas. For example, several products such as honey, meat and even beer have already been associated with the management of Natura 2000 site (Kettunen et al 2009¹⁶). Similarly, as the assessment in Section C shows, the network continues to provide significant opportunities for recreation and tourism, including related businesses. These revenue streams can provide a funding source that helps to cover some opportunity costs of Natura 2000 and also actively contribute towards the management of the sites (i.e. being channelled into covering the costs of management activities).

Public-private partnerships: The opportunities outlined above create possibilities for engaging a broader range of stakeholders in managing the Natura 2000 network. Several business sectors can be brought on-board, including companies depending on a steady supply of clean water, businesses benefiting from the natural beauty and other characteristics of the site, or businesses seeking to create markets for biodiversity-friendly products. The conservation and/or restoration of ecosystems and their services can provide cost-effective solutions for areas such as water management and flood mitigation, and as such are being increasingly considered as good investment by different businesses.

¹⁶ Kettunen, M., Bassi, S., Gantioler, S. & ten Brink, P. 2009. Assessing Socio-economic Benefits of Natura 2000 – a Toolkit for Practitioners (September 2009 Edition). Output of the European Commission project Financing Natura 2000: Cost estimate and benefits of Natura 2000 (Contract No.: 070307/2007/484403/MAR/B2). Institute for European Environmental Policy (IEEP), Brussels, Belgium. 191 pp. + Annexes.

Part E:

Summary of results and outlook

XII) Summary of results

The primary objective of the Natura 2000 protected area network is the conservation of Europe's unique and endangered biodiversity, including rare habitats, species and genetic diversity. In addition, the Natura 2000 network provides a wide range of benefits to society and the economy via the flow of ecosystem services (provisioning, regulating, cultural and supporting services). These also support policy objectives beyond biodiversity, in particular climate change mitigation and adaptation, water quality and provision, food provision, jobs and livelihoods, cost savings, science and education, social cohesion and identity.

Assessing the benefits associated with the Natura 2000 network is crucial, not only to underline the importance of conservation and the risks of degradation, but also to communicate the need for funding, help address stakeholders' (mis)perceptions of the importance of the sites, and help integrate the sites into the wider ecological-social-economic fabric of the regions. In addition, the EU has made commitments in Nagoya to the Strategic Plan of the Convention on Biological Diversity (CBD), which includes targets for the assessment of the benefits of natural capital and integration into national accounts. Further assessment of Natura 2000 benefits, and indeed those of wider green infrastructure and other living natural capital, will be essential for this



to be achieved and to ensure that policy makers at local, national and international level have the full evidence base available to take account of the value of Natura 2000 in their decisions.

Given existing data and available research, **this study derives some very broad estimates for the overall economic benefits of the Natura 2000 network across the EU.** This builds both on an analysis of existing site – and habitat-based valuations, as well as on an estimate for selected ecosystem services.

A first estimate for the value of the benefits of the (terrestrial) Natura 2000 network – scaling up from existing site-based studies – suggests that these could be **between €200 and €300 billion per year at present** (or 2% to 3% of EU GDP¹⁷). This value should be seen as ‘gross benefits’ delivered by sites, rather than the net benefits of the Natura 2000 designation and associated conservation measures.

The estimate is based on a relatively small number of studies scaled up to the EU level using the benefit transfer method. There are many methodological issues associated with this approach that are recognised in the report. This range should therefore be seen as an indicative first estimate, which would benefit from further refinement through subsequent analysis. Furthermore, it should be noted that this value includes both market and welfare values, so the comparison to GDP should be seen only as an illustration of scale.

The **ecosystem-service analysis** identified some preliminary values for a set of services. Among those selected for the assessment, it should be noted that some services were amenable to a relatively robust estimate (namely carbon storage and tourism) while others could only undergo a more illustrative/experimental analysis. Furthermore, the number of services covered is but a subset of the services that the network offers, hence offering only a partial picture of the total benefits of the network. Nevertheless, the mix of ecosystem services here estimated seems to broadly match with the overall site-based benefits assessment of €200 to €300 billion/year. Much more work, however, is clearly needed to derive a robust order-of-magnitude estimate (see next section) and the services contribution to the whole.

A summary of key results by ecosystem service is provided below. The individual estimated values, what they relate to, their level of ‘robustness’, and suggested future research needs is shown in Table 10.

Carbon: The Natura 2000 network provides a critically important service of storing carbon, revealing essential synergies of biodiversity with climate mitigation and adaption; improvements in land management will increase these carbon benefits

Thanks to the data and methodologies already available, the benefits of Natura 2000 associated with carbon storage are the most amenable to a quantitative and monetary assessment.

In general, Natura 2000 sites store carbon at relatively high densities compared to the EU land area as a whole. Many sites protect several ecosystems that are important

¹⁷ EU budget 2011 http://ec.europa.eu/budget/figures/2011/2011_en.cfm





current stores of carbon and offer significant opportunities for further carbon sequestration, including sites located on forested lands, wetlands (especially peatlands), agricultural lands, and marine and coastal ecosystems. It is estimated that the Natura 2000 network currently stores around 9.6 billion tonnes of carbon, equivalent to 35 billion tonnes of CO₂, which is estimated to be worth between €600 billion and €1,130 billion (stock value in 2010)¹⁸, depending on the price attached to a ton of carbon (i.e. to reflect the value of avoided damage of climate change by avoided GHG emissions).

It can be expected that in the future these carbon values will increase, especially if the conservation status of the network improves. A policy scenario (Policy ON), where full Protected Area coverage (terrestrial PAs + fuller MPAs) with a move to full favourable conservation status is estimated to generate a gain of at least a total of 1.71-2.86% by 2020 compared to a policy inaction scenario (Policy OFF), where no additional action is taken to conserve the current Natura 2000 sites over the next decade.

Natural hazards: Natura 2000 sites offer potential significant cost savings and reduction of damage from extreme weather events

Natural hazards cause significant damage across the EU. For the period 1990 – 2010, the value of economic losses from natural disasters in the EU-25 amounted to around €163 billion (or €16 billion a year). Moreover, due to demographic trends and impacts of climate change, it is likely that the vulnerability of human settlements to natural hazards will increase in the future.

Given the important functions that natural barriers and green infrastructures can provide, Natura 2000 sites can and have played a role in the mitigation of natural hazards, such as floods, avalanches or landslides. Using natural measures to mitigate impacts of natural disasters can lead to cost effective solutions which are often less expensive than manmade ones.

The site specific nature of natural hazards mitigation and the limited data availability on the role of Natura 2000 in reducing risks across Europe means that, at this stage, it is not yet possible to estimate Natura 2000 wide benefits.

¹⁸ To do: ensure that understandable for non-economists.

Tourism: Natura 2000 is already proving to be an important motor of many local economies by attracting tourists, whose spending supports local economies.

The expenditure supported by visitors to Natura 2000 sites is around €50-85 billion /year (in 2006). Only a share of the visitors is explicitly attracted by the Natura 2000 designation (most are simply attracted to the site for its aesthetic and landscape value). If only the expenditure of those visitors who have affinity for Natura 2000 designation is taken, the range becomes to €9-20 billion/year for 230-520,000 visitor days. The value of recreational benefits derived by the visitors themselves is estimated at €5-9 billion per annum.

Furthermore, protected areas can provide additional benefits to the **local and regional economy**, by attracting inward investment and enhancing local image and quality of life.

Marine Protected Areas: Marine Natura sites, as part of a wider network of connected marine areas, may have positive effects on overexploited fish stocks.

Marine Protected Areas (MPAs) support a range of ecosystem services.

The value of benefits delivered by the marine area currently protected by the network (equivalent to 4.7% of the EU's marine area) is approximately €1.4-1.5 billion per year. This would increase up to €3.0-3.2 billion per year if 10% of the sea area were protected, and €6.0-6.5 billion per year for protection of 20% of the sea area. The higher figures apply to stronger protection measures.

This should be seen as a ball park value, illustrative of the importance of this issue. Obtaining more robust results would need an improved understanding of how protection will influence habitats, services and off-site fisheries; the level to which benefits will depend on details of protection; and network effects.

Water: Money can be saved via working with natural capital, saving water purification and provisioning costs

While it has not been methodologically feasible to develop an EU wide assessment of the benefits of the Natura network for water purification and provision, given the site specific nature of the benefits, it is clear from case examples that the Natura 2000 network can lead to cost-effective means of water purification and supply, offering significant savings over man-made alternatives.

To cite an example from central and northern Europe, for the four European cities of Berlin, Vienna, Oslo and Munich, protected areas have been estimated to bring average per capita benefits ranging between €15 and €45 per year for both water purification and provision combined. This compares to average household water bills of €200 per year (in Germany). This underlines that benefits can be indeed significant, and lead to substantial actual and potential cost

savings from ecosystem based water purification and provision, both for companies (in terms of reduced operational costs) and citizens (reduced water bills). It will be important for cities to explore the role of natural capital (protected areas, wider green infrastructure) in the purification and provision of water and ensure that such considerations are integrated in the water management plans required under the Water Framework Directive.

Food security and provision:

Insect Pollination services are important in Europe – with an annual value estimated at €14 billion per year, which represents 10% of agricultural production for human food in 2005. However, the existing data does not allow identification of the share of this from Natura 2000 and from wider Green Infrastructure.

Many Natura 2000 sites also support important **agricultural practices**. Farmland covers almost 50% of the EU territory and agro-ecosystems represent 38% of surface of Natura 2000 sites. High Nature Value farming can offer significant benefits for biodiversity as well as helping to support local breeds, conserving genetic diversity and enhancing the resilience of the sector.

Table 10: Summary of results, what they relate to, the level of robustness and further needs in the area**Key:**

Deep green	Robust numbers – fine for publication, citation, without need for significant context.	Orange	Illustrative/indicative numbers but use with care and not out of context of the being a first assessment
Light green	Illustrative/indicative – useable with due caveats.	Red	Weak/very experimental. Do not use
Italics text	Experimental or illustrative	Bold text	Key point, result
Abbreviations:	Bn = billion;		yr = year

Approach	Numbers	What they relate to	Level of robustness/usability	Needs
Site Based	€ 223–314 bn/yr <i>€ 251–360 bn/yr</i>	Grossing up from 35 values from 21 studies. GDP adjusted site based Non adjusted site based	Best currently possible preliminary indicative value. Use with care; lot of caveats. High dependence on studies from UK and Netherlands.	To have robust order of magnitude ~ ideally a minimum of 200 comparable studies should be available – across biogeographic regions. A priority would be to get wider geographic focus.
Habitats Based	€ 189–308 bn/yr	Grossing up from 33 study numbers for 7 habitat groups – coastal, freshwater, heath and sand, grasslands, bogs and mires, forests.		A bottom up survey of ecosystem services (ESS) from sites and beneficiaries to help assess factors driving benefits.

Territorial (extrapolation from national based studies)	<i>n/a as rejected for this study</i>	<i>Grossing up from Scotland, E&W and NL to rest of the EU</i>	<i>Not robust/useable. Was useful as a Straw man in the study</i>	A possible way forward would be to focus on smaller territorial scale. A significant increase in stud- ies, noting biogeo- graphic regions as well as range of key site and context indicators.
Carbon sequestration/ storage	Current Stock: €600bn–€1130bn for the Natura 2000 network. Policy On: Over next 10 years, there will be an increase of €79-88 bn in carbon value if ecosystem qual- ity is improved; or €82-92 bn if there is a 10% increase in forest area.	Stock value from carbon storage (liv- ing + dead carbon) and CO ₂ value range €17.3 and 32/tCO ₂ for 2010. Annual sequestration build- ing on sequestration rates. Stock values are gross values. Policy-on values: increment.	Relatively robust estimate for the value of the stock of (living) carbon. Is an underestimate of the total value given that sequestration not addressed.	A further breakdown, site corroboration. More an annual natural gains – sequestration. Look also more at soil carbon given that this is a com- plex issue.
Natural hazards	<i>Context values:</i> €160 Bn over 1980 to 2010 i.e. ~€5bn year losses SR: country indica- tive estimate: €3.75 bn from restoration/planning	Country example: Slovak Republic: the national Landscape Revitalisation and Integrated River Basin Management Programme > ~ ben- efits of €3.75 bn, mainly related to avoiding the costs of flood protection measures (Gov't of Slovak Republic, 2010).	Unable to produce numbers related to Natura 2000. Note that the losses noted in the left column do not represent cur- rent costs avoided by Natura 2000 or green infrastructure. It is currently not possible to say what these would be.	Geographic Information Systems (GIS), survey of sites risks, impacts, role of Natura 2000, benefits.
Water – provision and purification	Provision Experimental ~ €22 billion/year Purification Experimental €2.2–€25 billion/ year	Provision Grossing up from 9 studies/values – but only 1 from the EU. Purification Grossing up from 3 values	EU values currently experimental; use case examples as these communi- cate the benefits	Survey/analy- sis of cities and beneficiaries.

Pollination	<i>Context values:</i> EU: Total € 14 bn/yr, which is 10% of agriculture productivity. World Pollination: € 153 bn/yr	General value of insect pollination	Order of magnitude robust, but not for Natura 2000 share. While it is clear that Natura 2000 sites are habitats for a wide range of wild pollinators for onsite agricultural activity and for nearby agricultural production, there is insufficient data to be able to allocate share to Natura.	Explore what the direct role of Natura 2000 is in wild pollination and the overall share. Useful to identify and assess specific sites that offer particular pollination value and output gains/input savings.
Marine	<i>Production services: food: fish</i> € 1 Bn per year off-site fisheries benefits <i>Wider set of Marine ESS:</i> € 2.5–3.8 Bn per year improvement in 7 services	Approximate marginal benefits associated with protecting 10% of EU marine environment; range is for less-more restrictive protection. Based on transfer of expert judgement.	<i>Highly uncertain, order of magnitude estimates. Fisheries value only ballpark, dependent on CFP reform.</i>	<i>Full habitat data (5 years); research and monitoring to understand the impacts of protection on services.</i>
Tourism expenditures	Around € 50–85 bn/yr (in 2006) for 1.7 billion visitor days (~466,000 visitors/day average) considering all visitors Between € 9–20 bn/yr considering visitors with affinity for Natura 2000 designation	Scaling up from a representative sample of 47 Natura 2000 sites	Order of magnitude rather than precise estimate (margin of error), comparable with economic indicators of tourism (e.g. the estimated value added of tourism and recreation for EU-27 is € 505 bn)	<i>More data on tourism at site level (number of visitors and tourism spending)</i> <i>Better determination of the affinity of visitors for Natura 2000 designation</i>
Recreation (non market benefits)	€ 4/visit i.e. between € 5–9 billion over the overall Natura 2000 network	Scaling up from a list of recreational values taken from the literature (National parks, Natura 2000 sites, habitats)	Rough order of magnitude rather than precise estimate, comparable with other recreational values for Natura 2000 sites	<i>More values from Natura 2000 case studies developed under a comparable protocol; values on activities and attractiveness of sites</i>

Overall, it should be acknowledged that there are a range of methods to ascertain value, and the values derived themselves can be of different types – from real market values that can feature in companies' 'bottom lines', national accounts and GDP, to values representing wellbeing, which are meaningful at a social level, but invisible to the cash economy. The values also accrue to a wide set of beneficiaries and will have very different implications for protected areas funding. Only a small proportion of the estimated benefits of €200–300 billion are reflected in cash transactions, and in reality very little actually accrues directly to protected areas. This underscores a fundamental issue: while protected areas have value to economies and societies, this value are generally not visible directly (hence the need for assessment) and their related benefits rarely pay the site manager. The protected areas are important public goods, creating many private benefits, but generally provide far less return for their ongoing management, maintenance or improvement of conservation status.

XIII) Outlook: **improving the knowledge base for the economic valuation of ecosystem services delivered by Natura 2000**

Currently, only a few ecosystem services can be quantified and valued for the Natura 2000 network as a whole, given limitations in data and methodology. However, with additional investment in data and studies, it is expected that a fuller and more robust assessment of the benefits of Natura 2000 network in the EU can be achieved in the next ten years.

Progress on economic valuation does not mean that other techniques (e.g. biophysical valuation, assessments or stakeholder assessments) become less relevant. On the contrary, **what is needed is progress with the range of tools to better define the contribution of nature to society and the economy as well as its intrinsic value.** Also, this study confirms that, while identifying the values of ecosystem services may be relatively feasible, measuring the ecosystem services specifically delivered by Natura 2000 sites remain a great challenge to economic valuation. This will be worth further efforts in future analysis.

It should be recognised that it will never be possible (nor, arguably, needed) to derive a precise, robust, static value for Natura 2000. The value will always be dynamic, affected by population growth, demography, income, changing geographic conditions, interests and preferences, economic contexts and wider contexts (e.g. global carbon values and climate change). This, and the site specificity of functions, services and values, also mean that there are limits to what can be assessed for the EU level as a whole.

Nevertheless, **there are strong merits in supporting the development of additional site-based benefits valuation for Natura 2000** in a manner that would allow a wider 'meta-analysis' to be carried out.

To allow a statistically significant (i.e., robust) analysis, **more data is clearly needed.** Ideally, data sources would at least encompass 200 quality comparable primary valuation studies on the benefits of Natura 2000 from across the EU Member States – i.e. around 20 studies per key factor driving benefits (standard rule of thumb to help get statistically significant answers). As temporal and spatial conditions are



important and methods evolving, some past studies will not be useable in the future and new studies will be needed, based on a common methodology that builds on Member States' and TEEB approaches. Realistically, in the future it will be possible to update only a few of the figures/services currently estimated (e.g. change carbon values used) and new evidence and figures will have to be developed using state of the art tools (e.g. building on MA, TEEB framework and advances in methods). This approach for valuation could for instance be incorporated into future LIFE Biodiversity projects as appropriate.

Primary valuation will also be crucial to expand the evidence base. Value evidence is important in determining what the benefits of different sites and management options are, and forms a key input to policy and decision processes, notably regarding designation, allowable activities, and management methods. This does not mean, however, that primary valuation is required for all sites. Because there are many similarities across sites, the services provided and the human populations benefiting from them, value transfer methods can be used.

The development of **'benefit production functions/value transfer functions'** would therefore be desirable for an EU wide assessment, in order to identify and characterise key factors driving the benefit values. In practice this should be done separately for terrestrial sites and for marine sites, given the quite different drivers of value. What is needed is a transparent framework enabling to compare analyses and build on results from different methods in different contexts. The Millennium Ecosystem Assessment (MA) and TEEB frameworks offer a useful basis for this.

The potential for advancing valuation through improved use of **Geographic Information Systems (GIS) and mapping** is also very significant. The progress of the work of the JRC and EEA in mapping and use of ecosystem service indicators underlines the scope here for important advances. This could help with site-based assessment as well as with wider regional assessments. It is expected that these tools will be of particular support to the future assessment of carbon storage and sequestration, and also for water supply, with potential even as regards pollination (see e.g. Figure 7) as well as flood control, which are currently very difficult to assess given site specificity.

It would also be valuable to do an **analysis/survey of the level of ecosystem service provision from the different sites to different stakeholders** (across geographic levels) to quantify the inter-connections and explore the quantitative scale of benefits – for example in relation to specific benefits from carbon storage and sequestration of specific sites, the number of cities and people benefiting from water provisioning and purification and natural hazard control, and so on.

Concluding remarks: Needs and concrete steps to realise the benefits from Natura 2000

There is a new evidence base that conserving and investing in our biodiversity makes sense for climate challenges, for saving money, for jobs, for food, water and physical security, for cultural identity, health, science and learning, and of course for biodiversity itself.

Full implementation of the actions included in the EU 2020 Biodiversity strategy will be essential to realise these wider economic benefits of Natura 2000. The results of the Commission's work on estimating the overall economic value of the benefits provided by the Natura 2000 network presented in this brochure demonstrate that investing in the designation, protection, management and monitoring of Natura 2000 makes also economic sense.

References

- ABPMer, Risk and Policy Analysts, & Jan Brooke Environmental Consultant Ltd. Arcadis Belgium, EFTEC and ECNC. 2011. Recognizing Natura 2000 benefits and demonstrating the Economic benefits of conservation measures – Development of a Tool for Valuing Conservation Measures.
- Balmford, A., Rodrigues, A. S. L., Walpole, M., ten Brink, P., Kettunen, M., Braat, L. and de Groot, R. (2008) *The Economics of Biodiversity and Ecosystems: Scoping the Science*, European Commission (Contract: ENV/070307/2007/486089/ETU/B2), Cambridge, http://ec.europa.eu/environment/nature/biodiversity/economics/pdf/scoping_science_report.pdf.
- Beare, D.J., Holker, F., Engelhard, G.H., McKenzie, E. and Reid, D. (2010) An unintended experiment in fisheries science: a war mediated protected area in the North Sea results in Mexican waves in fish numbers-at-age. *Naturwissenschaften*, 97(9): 797–808.
- BIO Intelligence Service, Ecotrans, OÄR and Dunira Strategy. 2011. *Estimating the Economic Value of the Benefits Provided by the Tourism/Recreation and Employment Supported by Natura 2000*.
- Bostedt, Göran and Leif Mattsson (1991), "Skogens betydelse för turismen: En samhällsekonomisk pilotst".
- Braat and ten Brink (eds) (2008) *The Cost of Policy Inaction – The case of not meeting the 2010 biodiversity target*. http://ec.europa.eu/environment/nature/biodiversity/economics/teeb_en.htm
- Brauman, K.A., Daily, G.C., Duarte, T.K., and Mooney, H.A. 2007. The nature and value of ecosystem services: an overview highlighting hydrologic services. *Annual Review of Environmental Resources* (32): 67–98.
- Brown, O., Crawford, A. and Hammill, A. (2006) *Natural Disasters and Resource Rights: Building Resilience, Rebuilding Lives*, International Institute for Sustainable Development, Manitoba, Canada.
- Butcher Partners Limited (2006). *Economic benefits of water in Te Papanui Conservation Park. Inception Report*.
- Campbell, A., Miles, L., Lysenko, I., Hughes, A., Gibbs, H. 2008. *Carbon storage in protected areas: Technical report*. UNEP World Conservation Monitoring Centre.
- Ceroni, M. (2007) *Ecosystem services and the local economy in Maramures Mountains Natural Park, Romania*. Unpublished report.
- Cooper, T., Hart, K. and Baldock, D. (2009) *The Provision of Public Goods Through Agriculture in the European Union*, Report Prepared for DG Agriculture and Rural Development, Contract No 30-CE-0233091/00-28, Institute for European Environmental Policy: London.
- Cruz, A de la, Benedicto, J., 2009. *Assessing Socio-economic Benefits of Natura 2000 – a Case Study on the ecosystem service provided by SPA PICO DA VARA/ RIBEIRA DO GUILHERME*. Output of the project Financing Natura 2000: Cost estimate and benefits of Natura 2000 (Contract No.: 070307/2007/484403/MAR/B2). 43pp http://ec.europa.eu/environment/nature/natura2000/financing/docs/azores_case_study.pdf
- Desmyttere, H. and Dries, L. (2002) *Natura 2000 Promoting the socio-economic benefits of Natura 2000. Case Study in the 'Pond Complex of Central-Limburg*.
- Dubgaard A, Kallesøe M F, Petersen M L and Ladenburg J (2002) *Cost-Benefit Analysis of the Skjern River Restoration Project*. Department of Economics and Natural Resources, Royal Veterinary and Agricultural University Copenhagen.
- Dudley, N. and Stolton, S. 2003. *Running Pure: The importance of forest protected areas to drinking water*. World Bank/WWF Alliance for forest conservation and sustainable use. WWF: Gland, Switzerland.
- Eagles, P. and Hillel, O. (2008) 'Improving protected area finance through tourism in protected areas', in SCBD (Secretariat of the Convention on Biological Diversity) *Protected Areas in Today's World: Their Values and Benefits for the Welfare of the Planet*, CBD Technical Series No 36, CBD, Montreal, <http://www.cbd.int/doc/publications/cbd-ts-36-en.pdf>, accessed 10 November 2009.
- Ebert et al., 2009. *Floodplain restoration along the lower Danube: A climate change adaptation case study*. *Climate and Development* 1 (2009) 212–219.
- Eftec (2008) *Wallasea Island Economic Benefits Study*. Final report submitted to East of England Development Agency.
- European Commission, (2005). *Assessment, monitoring and reporting of conservation status – Preparing the 2001-2007 report under Article 17 of the Habitats Directive (DocHab-04-03/03 rev.3)*. Note to the Habitat Committee. Brussels. <http://www.lcie.org/Docs/Legislation/DocHab-04-03-03%20rev3.pdf>
- European Commission. (2007). *European Flood Action Programme*. Available at http://ec.europa.eu/environment/water/flood_risk/consult.htm (last updated in November 2007).
- European Environment Agency (EEA) (2010) *Scaling up ecosystem benefits. A contribution to The Economics of Ecosystems and Biodiversity (TEEB) study*. EEA Report No 4/2010.

- European Environmental Agency (2004). Mapping the impacts of recent natural disasters and technological accidents in Europe. European Environment Agency, Copenhagen, Denmark. 48 pp.
- Everard, M. (2009) Using science to create a better place: ecosystem services case studies. Better regulation science programme. Environment Agency.
- Gallai, N., Salles, J.M., Settele, J., Vaissiere, B.E., (2009). Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecological Economics* 68 (3), 810–821.
- Gantioler S., Rayment M., Bassi S., Kettunen M., McConville A., Landgrebe R., Gerdes H., ten Brink P. (2010) Costs and Socio-Economic Benefits associated with the Natura 2000 Network. Final report to the European Commission, DG Environment on Contract ENV.B.2/SER/2008/0038. Institute for European Environmental Policy/GHK/Ecologic, Brussels 2010.
- GHK (2011) Benefits of Sites of Special Scientific Interest. Final report for Defra.
- Gren I-M, Groth K-H and Sylven M (1995) Economic Values of Danube Floodplains. *Journal of Environmental Management* (1995) 45, 333–345.
- Grizzetti B, Bouraoui F, Aloe A. (2007) Spatialised European Nutrient Balance. EUR 22692 EN. 2007. JRC36653.
- Haines-Young, R. and Potschin, M. (2009) 'The links between biodiversity, ecosystem services and human well-being', in D. Raffaelli and C. Frid (eds) *Ecosystem Ecology: A New Synthesis*, BES Ecological Reviews Series, Cambridge University Press, Cambridge.
- Hernandez S. and Sainteny G. 2008. Evaluation économique et institutionnelle du programme Natura 2000: étude de cas sur la plaine de la Crau. Lettre de la direction des études économiques et de l'évaluation environnementale. Hors Série N°08 – Juillet 2008. URL: http://www.natura2000.fr/IMG/pdf/CREDOC-D4E_4pages-natura2000_2008.pdf
- Hopkins J and Maxted N (2011) Crop Wild Relatives: Plant Conservation For Food Security. Natural England Research Report NER037. <http://publications.naturalengland.org.uk/publication/31031>
- Intergovernmental Panel on Climate Change, Climate Change (IPCC). (2001). Climate Change: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change. Edited by J.T. Houghton et al.
- IPCC (2000), Special Report on Land Use, Land Use Change and Forestry.
- ISDR – International Strategy for Disaster Reduction (2004) 'Living with Risk: A global review of disaster reduction initiatives', UN/ISDR, Geneva, Switzerland. URL: http://www.preventionweb.net/english/professional/publications/v.php?id=657&utm_source=pw_search&utm_medium=search&utm_campaign=search
- Jacobs (2004) Environment Group Research Report: An Economic Assessment of the Costs and Benefits of Natura 2000 Sites in Scotland, 2004 Final Report, The Scottish Government. 75 pp URL: <http://www.scotland.gov.uk/Resource/Doc/47251/0014580.pdf>
- Jacobs Scottish Executive Environment Group Research Report (2005), An Economic Assessment of the Costs and Benefits of Natura 2000 Sites in Scotland, Final Full report, Scottish executive and Scottish Natural Heritage 2005.
- JRC, (2011). A European Assessment Of The Provision Of Ecosystem Services: Towards An Atlas Of Ecosystem Services. Authors: Maes, J., Paracchini, M.L., Zulian, G. Joint Research Centre – Institute For Environment And Sustainability. EUR – Scientific And Technical Research Series – ISSN 1018-5593. ISBN 978-92-79-19663-8. DOI:10.2788/63557. Available at: <http://publications.jrc.ec.europa.eu/repository/handle/111111111/16103>
- Kaimowitz D. 2005. 'Forests and Human Health: Some Vital Connections'. Swedish CGIAR, Bogor, Indonesia.
- Kasina, J. M. et al., (2009). Economic benefit of crop pollination by bees: a case of Kakamega small-holder farming in western Kenya. *Journal of Economic Entomology*, 102(2), p.467-473. Available at: [Accessed August 1, 2011].
- Kazakova, Y., Pop, E. (2009). Assessing Socio-economic Benefits of Natura 2000 – a Case Study on the ecosystem services provided by Oaş-Gutâi Plateau and Igriş site, Maramures, Romania. Output of the project Financing Natura 2000: Cost estimate and benefits of Natura 2000 (Contract No: 070307/2007/484403/MAR/B2). 22 pp. + Annexes.
- Kettunen, M., Bassi, S., Gantioler, S. & ten Brink, P. 2009. Assessing Socio-economic Benefits of Natura 2000 – a Toolkit for Practitioners (September 2009 Edition). Output of the European Commission project Financing Natura 2000: Cost estimate and benefits of Natura 2000 (Contract No.: 070307/2007/484403/MAR/B2). Institute for European Environmental Policy (IEEP), Brussels, Belgium. 191 pp. + Annexes.
- Kettunen, M., Berghöfer, Bouamrane A M., Brunner, A., Chape S., Conner, N., Dudley, N., Ervin, J., Gidda, Morling P., . S. B., Mulongoy, K. J., Pabon, L., Seidl A., Stolton S., ten Brink P., Vakrou, A., (2011). Recognising the value of protected areas. In *TEEB in National Policy – The Economics of Ecosystems and Biodiversity in National and International Policy Making*. Edited by Patrick ten Brink, IEEP. Earthscan, London.

- Kettunen, M., Genovesi, P., Gollasch, S., Pagad, S., Starfinger, U. ten Brink, P. and Shine, C. 2008. Technical support to EU strategy on invasive species (IAS) – Assessment of the impacts of IAS in Europe and the EU (final module report for the European Commission). Institute for European Environmental Policy (IEEP), Brussels, Belgium. 43 pp. + Annexes. (draft version)
- Klein, A., Vaissière, B. E., Cane, J. H., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C., & Tscharntke, T. (2007). Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society of London B* 274: 303–313.
- Kniivila M, Ovaskainen V and Saastamoinen O (2002) Costs and benefits of forestconservation: regional and local comparisons in Eastern Finland. *J. Forest Economics* 8, 131–150 (2002).
- Kuik O, Brander L and Schaafsma M (2006) Globale Batenraming van Natura 2000 gebieden. IVM, Amsterdam.
- Kukreja, N. (2010). Organic Farming and Biodiversity in Europe: Examples from the Polar Circle to Mediterranean Regions. IFOAM EU Group. Brussels.
- Litskas, D., Mamolos, A., Kalburtji, K., Tsatsarelis, C. Kiose-Kampasakali, E. (2011). Energy flow and greenhouse gas emissions in organic and conventional sweet cherry orchards located in or close to Natura 2000 sites, *Biomass and Bioenergy*, Volume 35, Issue 3, March 2011, Pages 1302–1310, ISSN 0961-9534, 10.1016/j.biombioe.2010.12.023.
- Losey, J. E. and Vaughan, M. (2006) 'The economic value of ecological services provided by insects', *BioScience*, vol 56, pp311–323.
- Luisetti T, Turner R K, Hadley D and Morse-Jones S (2010) Coastal and marine ecosystem services valuation for policy and management. CSERGE Working Paper EDM 10-04.
- Maes, J., Paracchini, M.L., Zulian, G. 2011. A European assessment of the provision of ecosystem services. Towards and atlas of ecosystem services. Ispra: Joint Research Centre, IES.
- Maltby, E. (ed) (2009) *Functional Assessment of Wetlands: Towards Evaluation of Ecosystem Services*, Woodhead, Abington, Cambridge.
- Martin-Lopez B, Montes C and Benayas J (2007) Influence of user characteristics on valuation of ecosystem services in Donana Natural Protected Area (south-west Spain) *Environmental Conservation* 34 (3): 215–224.
- Millennium Ecosystem Assessment, 2005. *Ecosystems and Human Well-Being: Biodiversity Synthesis*. World Resources Institute, Washington, DC.
- Mücher, C.A., Hennekens, S.M., Bunce, R.G.H., Schaminee, J.H.J., and Schaepman, M.E. 2009. Modelling the spatial distribution of Natura 2000 habitats across Europe. *Landscape and Urban Planning* 92: 148–159.
- Mulongoy, K.J. and S.B. Gidda (2008). 'The Value of Nature: Ecological, Economic, Cultural and Social Benefits of Protected Areas', Secretariat of the Convention on Biological Diversity, Montreal, 30 pages.
- Pabian O., Jaroszewicz B. (2009) Assessing Socio-economic Benefits of Natura 2000 – a Case Study on the ecosystem service provided by Białowieża Forest. Output of the project Financing Natura 2000: Cost estimate and benefits of Natura 2000 (Contract No.: 070307/2007/484403/MAR/B2). 69pp. http://ec.europa.eu/environment/nature/natura2000/financing/docs/bialowieza_case_study.pdf
- Paracchini M.L., J.-E.Petersen, Y.Hoogeveen, C.Bamps, I.Burfield, C.van Swaay (2008): High Nature Value Farmland in Europe – An estimate of the distribution patterns on the basis of land cover and biodiversity data, Report EUR23480 EN. 87 p.
- Pearce, D.W. and Puroshothaman, S. (1992) *Protecting Biological Diversity: The Economic Value of Pharmaceutical Plants*. Discussion Paper 92-27. Centre for Social and Economic Research on the Global Environment, London.
- Penman, J., Gylarsky, M., Hiraishi, T., and T. Krug (eds.). 2003. *Good Practice Guidance for Land Use, Land Use Change, and Forestry*. Institute for Global Environmental Strategies for the Intergovernmental Panel on Climate Change. Available at <http://www.ipcc-nggip.iges.or.jp/public/gpoglulucf.htm> (17 August 2004)
- Pimentel, D. 2008. Conservation biological control. *Biological Control* 45: 171.
- Powe, Neil A. and Kenneth G. Willis (2002). *Mortality and Morbidity Benefits of Air Pollution Absorption by Woodland. Social & Environmental Benefits of Forestry Phase 2, Report To The Forestry Commission, Edinburgh*. Centre for Research in Environmental Appraisal and Management, University Of Newcastle Upon Tyne.
- Rensburg et al. (2009). *Socioeconomics of Farming for Conservation in the Burren*. BurrenLIFE Project.
- Ricketts, T. H., Daily, G. C., Ehrlich, P. R., & Michener, C. D. (2004). Economic value of tropical forest to coffee production. *Proceedings of the National Academy of Sciences* 101: 12579–12582.
- Ricketts, T. H., Regetz, J., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C., Bogdanski, A., Gemmill-Herren, B., Greenleaf, S. S., Klein, A. M., Mayfield, M. M., Morandin, L. A., Ochieng', A. and Viana, B. F. (2008), *Landscape effects on crop pollination*

- services: are there general patterns?. *Ecology Letters*, 11: 499–515. doi: 10.1111/j.1461-0248.2008.01157.x
- Roe D, Mulliken T, Milledge S, Mremi J, Mosha S and Grieg-Gran M. 2002. 'Making a Killing or Making a Living?', Wildlife trade, trade controls and rural livelihoods'. Biodiversity and Livelihood Issues number 6, IIED and TRAFFIC, London and Cambridge, UK.
- Ruijgrok E.C.M., Groot, R.S. de (2006). Kentallen Waardering Natuur, Water, Bodem En Landschap. Hulpmiddel Bij Mkb's. Available at http://www.mkbainderegio.nl/docs/Kentallen_waardering_natuur_water_bodem_en_landschap.pdf
- Ruijgrok, E.C.M. (2007) Blending ecology in actual economic decisions – the Dutch national guideline for ecosystem valuation applied on the Scheldt estuary in Belgium. Witteveen + Bos.
- Scialabba, N. (2003). Organic Agriculture: The challenge of sustaining food production while enhancing biodiversity. Sub-Group Meeting on Wildlife, Biodiversity and Organic Agriculture. Ankara (Turkey), 15-16 Apr 2003. FAO, Rome (Italy). Research, Extension and Training Div.
- Secretariat of the Convention on Biological Diversity (2010) Global Biodiversity Outlook 3. Montréal, 94 pages.
- Stolton, S.; Dudley, N.; Randall, J. (2008) Natural Security: Protected areas and hazard mitigation, WWF International, Gland, Switzerland.
- Sundseth, K. Undated. Natura 2000 and agriculture in Croatia. State Institute for nature protection, Croatia. www.natura2000.hr/UtilPages/GetDBDocument.aspx?id=47
- TEEB (2010) The Economics of Ecosystems and Biodiversity. Ecological and Economic Foundations. Edited by Pushpam Kumar. Earthscan, London and Washington.
- TEEB (2011) The Economics of Ecosystems and Biodiversity in National and International Policy Making. Edited by Patrick Ten Brink, Earthscan, London and Washington.
- TEEBcase by J. Förster (2010) Peatlands restoration for carbon sequestration, Germany, available at: [TEEBweb.org; http://www.eea.europa.eu/atlas/teeb/peatland-restoration-for-carbon-sequestration-germany-1](http://www.eea.europa.eu/atlas/teeb/peatland-restoration-for-carbon-sequestration-germany-1)
- UK National Ecosystem Assessment (2011) The UK National Ecosystem Assessment. UNEP-WCMC, Cambridge.
- Van der Ploeg, S. and R.S. de Groot (2010) The TEEB Valuation Database – a searchable database of 1310 estimates of monetary values of ecosystem services. Foundation for Sustainable Development, Wageningen, The Netherlands.
- Vienna Waterworks. 2011. General water quality in Vienna. Available at: <http://www.wien.gv.at/english/environment/watersupply/quality/quality-general.html>
- Willis, K.G (1990) Valuing non-market wildlife commodities: an evaluation and comparison of benefits and costs. *Applied Economics*. Volume 22, Issue 1, 1990, Pages 13 – 30.

Annex I: Glossary of terms¹⁹

Altruistic value: The importance which individuals attach to a resource that can be used by others in the current generation, reflecting selfless concern for the welfare of others (intra-generational equity concerns).

Avoided Costs: The costs that would have been incurred in the absence of ecosystem services.

Benefits: positive change in wellbeing from the fulfilment of needs and wants.

Bequest value: The importance individuals attach to a resource that can be passed on to future generations, reflecting intergenerational equity concerns.

Biodiversity: the variability among living organisms, including terrestrial, marine, and other aquatic ecosystems. Biodiversity includes diversity within species, between species, and between ecosystems (UN, 1993).

Biome: a large geographic region, characterized by life forms that develop in response to relatively uniform climatic conditions. Examples are tropical rain forest, savannah, desert, tundra.

Biophysical valuation: A method that derives values from measurements of the physical costs (e.g., in terms of labour, surface requirements, energy or material inputs) of producing a given good or service.

Carbon sequestration: The process of increasing the carbon content of a reservoir other than the atmosphere. (MA, 2005)

Consumer surplus: The benefits enjoyed by consumers as a result of being able to purchase a product for a price that is less than the most that they would be willing to pay.

Contingent valuation: Stated preference based economic valuation technique based on a survey of how much respondents would be willing to pay for specified benefits. (MA, 2005)

Cost-benefit analysis: A technique designed to determine the feasibility of a project or plan by quantifying its costs and benefits. (MA, 2005)

Direct use value (of ecosystems): The benefits derived from the services provided by an ecosystem that are used directly by an economic agent. These include consumptive uses (e.g., harvesting goods) and non-consumptive uses

(e.g., enjoyment of scenic beauty). Agents are often physically present in an ecosystem to receive direct use value. (MA, 2005)

Driver (direct or indirect): any natural or human-induced factor that directly or indirectly causes a change in an ecosystem.

Ecosystem: Means 'a dynamic complex of plant, animal and micro-organism communities and their non-living environment interacting as a functional unit' (UN, 1993).

Ecosystem degradation: A persistent reduction in the capacity to provide ecosystem services. (MA, 2005)

Ecosystem services: the direct and indirect contributions of ecosystems to human well-being. The concept 'ecosystem goods and services' is synonymous with ecosystem services.

Ecotourism: Travel undertaken to access sites or regions of unique natural or ecologic quality, or the provision of services to facilitate such travel.

Existence value: the value that individuals place on knowing that a resource exists, even if they never use that resource (also sometimes known as conservation value or passive use value).

Favourable Conservation Status: In layman's terms can be described as a situation where a habitat type or species is prospering (in both quality and extent/population) and with good prospects to do so in future as well²⁰.

Hedonic pricing: An economic valuation approach that utilizes information about the implicit demand for an environmental attribute of marketed commodities.

Human well-being: concept prominently used in the Millennium Ecosystem Assessment – it describes elements largely agreed to constitute 'a good life', including basic material goods, freedom and choice, health and bodily well-being, good social relations, security, peace of mind, and spiritual experience.

Incentives (disincentives), economic: a material reward (or punishment) in return for acting in a particular way which is beneficial (or harmful) to a set goal.

Indirect-use value (of ecosystems): the benefits derived from the goods and services provided by an ecosystem that are used indirectly by an economic agent. For example, the purification of drinking water filtered by soils.

¹⁹ Building on TEEB (2010) and TEEB (2011)

²⁰ Assessment, monitoring and reporting of conservation status – Preparing the 2001-2007 report under Article 17 of the Habitats Directive (DocHab-04-03/03 rev 3)

Intrinsic value: The value of someone or something in and for itself, irrespective of its utility for someone else. (MA, 200a)

Natural capital: an economic metaphor for the limited stocks of physical and biological resources found on earth, and of the limited capacity of ecosystems to provide ecosystem services.

Non-use value: benefits which do not arise from direct or indirect use.

Opportunity costs: foregone benefits of not using land/ecosystems in a different way, e.g. the potential income from agriculture when conserving a forest.

Primary valuation studies: Empirical valuation studies rather than those that rely on the transfer of values or value functions from other studies.

Production function: A function used to estimate how much a given ecosystem service (e.g., regulating service) contributes to the delivery of another service or commodity which is traded on an existing market.

Public goods: a good or service in which the benefit received by any one party does not diminish the availability of the benefits to others, and where access to the good cannot be restricted.

Replacement cost: The costs incurred by replacing ecosystem services with artificial technologies.

Resilience (of ecosystems): their ability to function and provide critical ecosystem services under changing conditions.

Revealed preference: A method to assess possible value options or to define utility (consumer preferences) based on the observation of consumer behaviour.

Scale: The measurable dimensions of phenomena or observations. Expressed in physical units, such as meters, years, population size, or quantities moved or exchanged. In observation, scale determines the relative fineness and coarseness of different detail and the selectivity among patterns these data may form. (MA, 2005)

Stakeholder: A person, group or organization that has a stake in the outcome of a particular activity.

Stated preference: Consumer preferences are understood through questions regarding willingness to pay or willingness to accept.

Substitutability: The extent to which human made capital can be substituted for natural capital (or vice versa).

Threshold/tipping point: a point or level at which ecosystems change, sometimes irreversibly, to a significantly different state, seriously affecting their capacity to deliver certain ecosystem services.

Total economic value (TEV): a framework for considering various constituents of value, including direct use value, indirect use value, option value, quasi-option value, and existence value.

Trade-offs: a choice that involves losing one quality or service (of an ecosystem) in return for gaining another quality or service. Many decisions affecting ecosystems involve trade-offs, sometimes mainly in the long term.

Travel cost method: A revealed preference valuation method that infers the value of a change in the quality or quantity of a recreational site (e.g., resulting from changes in biodiversity) from estimating the demand function for visiting the site.

Valuation: The process of expressing a value for a particular good or service in a certain context (e.g., of decision-making) usually in terms of something that can be counted, often money, but also through methods and measures from other disciplines (sociology, ecology, and so on). (MA, 2005)

Valuation, economic: the process of estimating a value for a particular good or service in a certain context in monetary terms.

Value: The contribution of an action or object to user-specified goals, objectives, or conditions. (MA, 2005)

Vulnerability: Exposure to contingencies and stress, and the difficulty in coping with them.

Willingness to pay: The maximum amount that a person is willing to pay for a good they do not have.

Willingness-to-pay (WTP): estimate of the amount people are prepared to pay in exchange for a certain state or good for which there is normally no market price (e.g. WTP for protection of an endangered species).

Annex II: Overview of existing valuation site-based studies

All values have been estimated on a per hectare per annum basis, where necessary by estimating the annualised values where the source material expressed these as capitalised sums, and have been converted to euro at 2011 prices.

The per hectare values are derived from estimates of the value of services delivered by each site, divided by the area of the site. It is apparent from Table A.2 that the available estimates give a wide range of values for the benefits of Natura 2000 sites, ranging from just less than €50 per hectare per year to almost €20,000 per hectare per year.

Table A.2: Summary of valuation studies, by site

Site	Ecosystem services/types of benefit	Site value per ha per year (€, 2011 prices)	Reference
Pond Complex of Central-Limburg, Belgium	Provisioning services, tourism and recreation.	1,406	Desmyttere and Dries (2002)
Scheldt estuary, Belgium	Regulating and provisioning ES (various).	3,990	Ruijgrok, E.C.M. (2007)
Skjern River restoration, Denmark	Biodiversity/existence values, recreation, water purification and regulation, fibre production.	1,218	Dubgaard et al (2002)
Protected forests in eastern Finland	Non market values measured through contingent valuation.	403	Kniivila et al (2002)
La Crau, France	Non-market benefits (public WTP) + hay production.	229	Hernandez and Sainteny (2008)
Donana, Spain	Range of ecosystem services, estimated through CVM.	375	Martin-Lopez et al (2007)
Sites protected for Large Blue butterfly, Landau, Germany	Range of services and values including non-use values.	6,932	Wätzold et al. (2008)
Burren, Ireland	Cultural services: tourism and recreation; Broader socio-economic benefits: beneficial externalities of conservation.	2,714	Rensburg et al. (2009)

Wadden Sea N2K sites, Netherlands	Wide range of provisioning, regulating and cultural services.	3,650	Kuik et al (2006)
River N2K sites, Netherlands	Use and non use values, estimated through hedonic pricing and benefit transfer.	5,324	
Lake and marsh N2k sites, Netherlands	Tourism, recreation, non use values including biodiversity	5,944	
Dune N2K sites, Netherlands	Flood protection, recreation, non use values.	13,198	
High fen and sandy soil N2K sites, Netherlands	Recreation, non use values.	1,274	
Stream valley and hills N2K sites, Netherlands	Provisioning, amenity, recreation, non-use values measured through stated and revealed preference methods.	4,974	
Białowieża Forest, Poland	Recreation, amenity and existence, freshwater, range of provisioning services (e.g. food, timber), tourism, pest control.	2,799	Pabian and Jaroszewicz (2009)
Pico da Vara/Ribeira do Guilherme, Azores, Portugal	Water provision, quality & regulation. Recreation and eco-tourism. Landscape and amenity values.	642	Cruz and Benedicto (2009)
Lower Green Corridor, Romania	Provisioning services: fisheries, forestry, animal fodder; Regulating services: nutrient retention; Cultural services: recreation.	512	Ebert et al. (2009)
Danube floodplains (7 countries, 60% in Romania)	Provisioning services, recreation, water purification.	572	Gren et al (1995)
Maramures Mountains Natural Park, Romania	All ecosystem services.	416	Ceroni (2007)

Clyde Valley Woods, Scotland	Recreation and non-use values (based on CVM of visitors and general public.	5,665	Jacobs (2004)
Waukenwae and Red Mosse, Scotland	Recreation and non-use values (based on CVM of visitors and general public.	14,769	
River Bladnoch, Scotland	Recreation and non-use values (based on CVM of visitors and general public.	5,341	
Sands of Forvie, Scotland	Recreation and non-use values (based on CVM of visitors and general public.	4,404	
Tips of Corsemaul and Tom Mor, Scotland	Recreation and non-use values (based on CVM of visitors and general public.	19,763	
Strathglass Complex, Scotland	Recreation and non-use values (based on CVM of visitors and general public.	87	
Lewis and Harris, Scotland	Recreation and non-use values (based on CVM of visitors and general public.	155	
Sites of special scientific interest in England and Wales (almost 80% by area are N2K)	Range of 7 key provisioning, regulating and cultural services (gross).	7,926	GHK (2011)
Wallasea Island, England	Range of key ecosystem services.	1,447	Eftec (2008)
Derwent Ings , England	Social benefits of N2K site, measured through CVM.	1,318	Willis, K.G (1990)
Skipworth Common, England	Social benefits of N2K site, measured through CVM.	5,987	
Upper Teasdale, England	Social benefits of N2K site, measured through CVM.	1,150	
Alkborough Flats, North Lincolnshire, England	Range of ecosystem services.	4,508	Everard, M. (2009)
Humber Estuary, England	Amenity and recreation, carbon.	847	Luisetti et al (2010)
Blackwater Estuary, England	Amenity and recreation, carbon, fisheries.	4,371	

Anex III: Examples of ecosystem services

The Table below shows a brief description of the different ecosystem services and provides some illustrative examples.

Table A3: Examples of ecosystem services

Ecosystem service	Ecosystem Service description	Illustrative example
Provisioning Services		
Food	Natura 2000 can play a significant role by providing fish, directly supporting sustainable agricultural production, such as through organic farming, and indirectly supporting out-of-the-site agricultural production (i.e. through wild pollination, erosion control, water cycling etc.). Moreover, some Natura 2000 sites also provide various wild products, such as mushrooms, berries or game.	Being the first major farming for conservation project in Ireland, The BurrenLIFE Project seems to offer a good 'value-for-money' solution with minimum estimated economic return of 235%. (Rensburg et al., 2009)
Water quantity	Ecosystems play a vital role in the global hydrological cycle, as they regulate the flow of water. Vegetation and forests influence the quantity of water available locally.	The benefits of freshwater provided by the Pico da Vara/Ribeira do Guilherme Natura 2000 park in Portugal are valued approximately € 600,000 per year or € 99 per hectare. Cruz and Benedicto (2009)
Raw materials	Ecosystems provide a great diversity of raw materials needed for instance for construction and fuel including wood, biofuels and plant oils that are directly derived from wild and cultivated plant species. There are also important Ornamental resources – Sustainably produced/harvested ornamental wild plants, wood for handcraft, seashells etc. Also ornamental fish.	Non-timber forest products such as rubber, latex, rattan and plant oils are very important in trade and subsistence – the annual global trade in such products is estimated to amount to US\$11 billion (Roe et al. 2002).
Natural medicines – Biochemicals & pharmaceuticals	Biodiverse ecosystems provide many plants used as traditional medicines as well as providing raw materials for the pharmaceutical industry. All ecosystems are a potential source of medicinal resources.	80% of the world's people are still dependent on traditional herbal medicine (WHO 2002), while the sale of medicines derived from natural materials amounts to US\$57 billion per year (Kaimowitz 2005).

Genetic/species diversity maintenance	Genetic diversity (the variety of genes between, and within, species populations) distinguishes different breeds or races from each other, providing the basis for locally well-adapted cultivars and a gene pool for developing commercial crops and livestock. Some habitats have an exceptionally high number of species which makes them more genetically diverse than others and are known as 'biodiversity hotspots'. In Europe, Mediterranean Basin with its particularly diverse flora is considered such a hotspot.	Crop Wild Relatives (CWR) are the wild ancestors of crop plants and other species closely related to crops. Hopkins and Maxted (2011) observed that they are likely to play a significant role in securing 21st century food security, because of their potential use in plant breeding to produce crops which withstand adverse impacts of climate change, increasing scarcity of nutrients, water and other inputs, and new pests and diseases.
Regulating services		
Air quality regulation	Trees or other plants also play an important role in regulating air quality by removing pollutants from the atmosphere. Many protected areas located in proximity to highly polluted areas might offer particularly high benefits.	The results of a study (Powe, 2002) have found net pollution absorption by trees in the UK to have reduced the number of deaths brought forward by air pollution by between 65-89 deaths and between 45-62 hospital admissions, with the net reduction in costs estimated to range somewhere between £222,308 and £11,213,276.
Climate/climate change regulation	Ecosystems regulate the global climate by storing and sequestering greenhouse gases. As trees and plants grow, they remove carbon dioxide from the atmosphere and effectively lock it away in their tissues. In this way forest ecosystems are carbon stores. Trees also provide shade whilst forests influence rainfall and water availability both locally and regionally.	In Mecklenburg-Vorpommern (Germany) an area of 29,764 ha (equivalent to about 10% of the area of drained peatlands in Mecklenburg-Vorpommern), has been restored between 2000 and 2008. This means that emissions of about 300,000 tCO ₂ -equivalents every year are avoided (with an average of 10.4 tCO ₂ -equivalents per hectare). When assuming a marginal cost of damage caused by carbon emissions of €70 per tCO ₂ , the effort to restore peatlands avoids damage from carbon emissions of up to €21.7 million every year, on average €728 per hectare of restored peatlands. (TEEB Case study by Förster 2011 and the references within ²¹)

²¹ <http://www.eea.europa.eu/atlas/teeb/peatland-restoration-for-carbon-sequestration-germany-1>

Moderation of extreme events	Ecosystems and living organisms create buffers against natural disasters, thereby preventing or reducing damage from extreme weather events or natural hazards including floods, storms, avalanches and landslides.	In the Swiss Alps, healthy forests are a major component of disaster prevention. 17 of Swiss forests are managed to protect against avalanches, landslides and rock falls. These services are valued at €1.6–2.8 billion per year (ISDR, 2004, Dudley et al., 2009).
Water regulation	Certain ecosystems, such as wetlands or sand dunes, can influence the timing and magnitude of water runoff, regulate and mitigate floods and provide support to recharging of ground water resources.	In Kalkense Meersen Natura 2000 site, in Belgium, it has been estimated that restoration of the original river landscape can bring flood mitigation benefits between €640,000–1,654,286 per annum (Arcadis Belgium et al., 2011).
Water purification & waste management	Ecosystems play a vital role in providing numerous cities with drinking water, as they ensure the flow, storage and purification of water. Furthermore, ecosystems such as wetlands filter effluents. Through the biological activity of microorganisms in the soil, most waste is broken down. Thereby pathogens (disease causing microbes) are eliminated, and the level of nutrients and pollution is reduced.	The city of Vienna obtains almost all of its drinking water from mountain springs originating in the Lower Austrian-Styrian high alpine zones. In December 2001, it was the first city in the world to protect its drinking water for future generations under Constitutional Law (Vienna Waterworks 2011).
Erosion control	Soil erosion is a key factor in the process of land degradation, desertification and hydroelectric capacity. Vegetation cover provides a vital regulating service by preventing soil erosion. Soil fertility is essential for plant growth and agriculture and well-functioning ecosystems supply soil with nutrients required to support plant growth.	A study by Ruijgrok et al. (2006) estimated that the value of erosion control in pristine scrubland areas in Europe and in Belgian grasslands was €44.5/ha, at 2008 prices (as in Braat et al, 2008).
Pollination	Insects and wind pollinate plants which is essential for the development of fruits, vegetables and seeds. Animal pollination is an ecosystem service mainly provided by insects but also by some birds and bats. Protected areas play a key role in harbouring wild pollinators which, if located in close proximity to agricultural fields, can help to increase yield and quality of many crops.	Using the methods of Gallai et al. (2009), the United Kingdom's National Ecosystem Assessment estimated the economic value of biotic pollination as a contribution to crop market value in 2007 at €629 million (England: €532 million, Northern Ireland: €28 million, Scotland: €69 million, Wales: unknown) (UK NEA, 2011)

Biological control	Ecosystems are important for regulating pests and vector borne diseases that attack plants, animals and people. Healthy ecosystems can effectively regulate pests and diseases through the activities of predators and parasites. Birds, bats, flies, wasps, frogs and fungi all act as natural controls.	Globally, more than 40 of food production is being lost to insect pests, plant pathogens, and weeds, despite the application of more than 3 billion kilograms of pesticides to crops, plus other means of control (Pimentel 2008).
Disease regulation of human health Regulation of vectors for pathogens	A number of species, such as birds and insects, are known to be vectors of human diseases (e.g. malaria, dengue fever, Lyme disease etc.). In a natural state the functioning of ecosystems keeps the populations of these species under control.	Asian tiger mosquito (<i>Aedes albopictus</i>) in Italy poses a health risk as it is a vector for Dengue and Chikunguna fever and it also has painful stings. Costs related to preventing negative health impacts (e.g. eradication program and communication) amounts to € 1.1 million/year (Kettunen et al. 2008 and the sources within).
Cultural & social services		
Landscape & amenity values	People around the world derive aesthetic pleasure from natural over built environment. In particular, people value a specific or exceptional view (landscape values) and appreciate the beauty of nature (amenity values).	'In Denmark, houses in natural environments, when compared to similar houses elsewhere, sell for a 25 percent higher price (Dissing, 2002). This is particularly true where they are located within 30-45 minutes of an urban centre (e.g. Danish Lille Vildmose site) (Bostedt et al., 1991).'
Ecotourism & recreation	Ecosystems and biodiversity play an important role for many kinds of tourism which in turn provides considerable economic benefits and is a vital source of income for many countries. Cultural and eco-tourism can also educate people about the importance of biological diversity. Walking and playing sports in green space is a good form of physical exercise and helps people to relax.	'Non-market benefits of the Scottish Natura 2000 sites related to recreation were estimated by asking visitors how much they would be willing to pay for using the Natura 2000 sites for recreational activities which resulted in an estimate of around £1.5 million per year related to use values. (Jacobs report to Scottish Executive, 2005)'
Cultural values and inspirational services, e.g. education, art and research	Language, knowledge and the natural environment have been intimately related throughout human history. Biodiversity, ecosystems and natural landscapes have been the source of inspiration for much of our art, culture and increasingly for science.	'The Bialowieza Forest, a Natura 2000 site, is the focus of extensive scientific research. Bialowieza village has three scientific institutes and two education centres. The national park runs a Museum and Bison Reserve with highly educated staff and a good level of nature education on offer.' Pabian and Jaroszewicz (2009)

Sources: Building on TEEB 2011b, TEEB 2010, MA 2005; Kettunen et al 2009; Balmford et al 2008; TEEB Foundations 2010a

European Commission

The Economic benefits of the Natura 2000 Network

Luxembourg: Publications Office of the European Union

2013 – 74 pp. – 21 x 29.7 cm

ISBN 978-92-79-27588-3

doi:10.2779/41957

ISBN 978-92-79-27588-3



Publications Office